

Archean to Recent aeolian sand systems and their sedimentary record: Current understanding and future prospects

JUAN PEDRO RODRÍGUEZ-LÓPEZ*, LARS B. CLEMMENSEN†, NICK LANCASTER‡, NIGEL P. MOUNTNEY§ and GONZALO D. VEIGA¶

**Departamento de Estratigrafía, Facultad de Ciencias Geológicas, Universidad Complutense de Madrid, Ciudad Universitaria, 28040 Madrid, Spain (E-mail: jprodrig@geo.ucm.es)*

†*Department of Geosciences and Natural Resource Management, Section of Geology, University of Copenhagen, Øster Voldgade 10, 1350 København K, Denmark*

‡*Division of Earth and Ecosystem Sciences, Desert Research Institute (DRI), 2215 Raggio Parkway, Reno, NV 89512, USA*

§*School of Earth and Environment, University of Leeds, Leeds LS2 9JT, UK*

¶*Centro de Investigaciones Geológicas, Universidad Nacional de La Plata-CONICET, Calle 1 # 644, B1900TAC La Plata, Argentina*

Associate Editor – Charlie Bristow

ABSTRACT

The sedimentary record of aeolian sand systems extends from the Archean to the Quaternary, yet current understanding of aeolian sedimentary processes and product remains limited. Most preserved aeolian successions represent inland sand-sea or dunefield (erg) deposits, whereas coastal systems are primarily known from the Cenozoic. The complexity of aeolian sedimentary processes and facies variability are under-represented and excessively simplified in current facies models, which are not sufficiently refined to reliably account for the complexity inherent in bedform morphology and migratory behaviour, and therefore cannot be used to consistently account for and predict the nature of the preserved sedimentary record in terms of formative processes. Archean and Neoproterozoic aeolian successions remain poorly constrained. Palaeozoic ergs developed and accumulated in relation to the palaeogeographical location of land masses and desert belts. During the Triassic, widespread desert conditions prevailed across much of Europe. During the Jurassic, extensive ergs developed in North America and gave rise to anomalously thick aeolian successions. Cretaceous aeolian successions are widespread in South America, Africa, Asia, and locally in Europe (Spain) and the USA. Several Eocene to Pliocene successions represent the direct precursors to the present-day systems. Quaternary systems include major sand seas (ergs) in low-latitude and mid-latitude arid regions, Pleistocene carbonate and Holocene–Modern siliciclastic coastal systems. The sedimentary record of most modern aeolian systems remains largely unknown. The majority of palaeoenvironmental reconstructions of aeolian systems envisage transverse dunes, whereas successions representing linear and star dunes remain under-recognized. Research questions that remain to be answered include: (i) what factors control the preservation potential of different types of aeolian bedforms and what are the characteristics of the deposits of different bedform types that can be used for effective reconstruction of original bedform morphology; (ii) what specific set of controlling conditions allow for sustained bedform climb versus episodic sequence accumulation and preservation; (iii) can sophisticated four-dimensional models be developed for complex patterns of spatial and

temporal transition between different mechanisms of accumulation and preservation; and (iv) is it reasonable to assume that the deposits of preserved aeolian successions necessarily represent an unbiased record of the conditions that prevailed during episodes of Earth history when large-scale aeolian systems were active, or has the evidence to support the existence of other major desert basins been lost for many periods throughout Earth history?

Keywords Aeolian, Archean, dunes, ergs, Mesozoic, Neogene, Palaeogene, Palaeozoic, preservation, Proterozoic, Quaternary.

INTRODUCTION

How can geologists best account for the preserved expression of aeolian sedimentary successions and relate such deposits to the varied set of processes responsible for their generation? “*The answer my friend is blowin’ in the wind*” (Dylan, 1963), and has been for at least 3.2 billion years. The aim of this study is to present an overview of the current state of the science relating to the sedimentology of aeolian sand systems and their preserved successions. Specific objectives are as follows: (i) to demonstrate the variability and complexity of the sedimentology of recent and ancient aeolian sand systems; (ii) to show how the spatial and temporal distribution of aeolian systems and preserved successions has varied throughout Earth history; (iii) to discuss the main mechanisms for the construction, accumulation and preservation of aeolian systems; and (iv) to present some future perspectives relating to issues that currently remain unresolved in aeolian sedimentology, thereby highlighting research targets and opportunities for the future. This study is supported by a suite of complementary material arranged in a series of tables that detail many of the best-known and most representative examples of siliciclastic, as well as some carbonate aeolian sand seas and coastal dunefields from the Archean and Proterozoic, Palaeozoic, Mesozoic and Cenozoic eras (see also Blakey *et al.*, 1988; Tedford *et al.*, 2005; Veiga *et al.*, 2011a; Simpson *et al.*, 2012). Although this work represents an attempt to compile an authoritative database of case-study examples for all periods in Earth history, many smaller and lesser-known aeolian systems have been omitted due to space limitations. The references contained in the supplementary tables of case studies (together with those references cited in the main manuscript) are contained in the supplementary file entitled ‘References text and tables’.

AEOLIAN SAND SYSTEMS AND THEIR SEDIMENTARY RECORD: CURRENT UNDERSTANDING

Aeolian sand systems can be divided into inland sand sea and coastal dune systems. Inland aeolian sand seas (also known as ergs) and the aeolian dunefields present within these large-scale sediment systems comprise bedforms of different morphological types and sizes (ranging from ripples to megadunes or draas), areas of sand sheets, interdunes (including non-aeolian sediments), as well as related extradune environments of alluvial, fluvial, lacustrine and marine affinity. Coastal dunefields likewise comprise various aeolian bedforms; many of these dune types – such as parabolic dunes – are also seen in inland systems, whereas others – such as coast-parallel dune ridges – are unique to coastal systems. Associated sediments include beach, wash-over fan and lagoon facies.

Following Kocurek (1999), the creation of an aeolian stratigraphic record can be considered in three phases (Fig. 1): (i) sand-sea (i.e. dunefield) construction; (ii) aeolian accumulation; and (iii) preservation of that accumulation. Construction of the world’s largest modern, active sand seas occurs in arid regions that typically experience less than 150 mm annual precipitation, although sites of significant aeolian construction also occur in non-desert settings, especially along sandy coastlines. Although aeolian sediment transport takes place under a wide range of wind energy regimes (Fryberger, 1979), it is the directional variability of such regimes that plays a major role in determining dune type, therefore dictating the range of sedimentary structures that develop on bedforms, and the style and rate of accumulation of deposits of those bedforms (Wasson & Hyde, 1983). Many present-day actively constructing and accumulating sand seas are located at sites of relatively lower wind energy compared with upwind areas, such that




Fig. 1. Schematic diagram showing the three-phase creation of the aeolian rock record and main controlling factors as proposed by Kocurek (1999).

sediment transport rates tend to decrease in the direction of transport, thereby encouraging sand deposition and accumulation. A down-wind reduction in sediment transport rate that leads to aeolian construction and accumulation may result from regional changes in atmospheric circulation patterns and wind regimes, whereby wind speed decreases and/or directional variability increases (Wilson, 1973; Lancaster, 1999, 2013). Coastal dunefields typically develop along lowland coasts where plentiful sediment supply (often beach sand) is available for inland transport by persistent onshore winds (e.g. Klijn, 1990). The size and morphology of coastal dunes are dependent on vegetation cover, sand supply, beach-dune interaction, wind regime and coastal orientation with regard to persistent winds.

Aeolian dunefield construction (the initiation and growth of systems of bedforms) is a function of sediment supply, the availability of that supply for aeolian transport and the transport capacity of the wind (Kocurek & Lancaster, 1999). Sediment supply is the volume of sediment suitable for aeolian transport generated per unit time; supply may be contemporaneous or time-lagged (Kocurek, 1999) and can be derived from multiple sources. The proximity of a dunefield to its sediment source area is reflected in the response of the system to changes in boundary conditions. Dunefields that lie close to their sediment source (including most coastal dunefields) tend to be sensitive to variations in sediment supply, whereas systems that develop far from their ultimate source tend to be more

sensitive to changes in sediment mobility or availability. Many sand seas are the depositional sinks of local to regional-scale sediment transport systems. Mineralogical and geochemical studies, aided in some instances by remote sensing data, can establish clear relations between source areas and sediment sinks (e.g. Scheidt *et al.*, 2011; Garzanti *et al.*, 2012). In many areas, however, these relations are not clear, and the source(s) of sand for major sand seas in the Sahara and elsewhere is (are) poorly constrained (Garzanti *et al.*, 2003; Muhs, 2004). Regional wind patterns appear to show long-distance transport paths in the Sahara and Australia, but recent work also points to the importance of local sources in Australia (Pell *et al.*, 2000) and elsewhere (Muhs *et al.*, 2003). The sand in coastal dunefields is derived primarily from the beach; textural and geochemical studies of foredune deposits can give information on sediment provinces and transport pathways in the nearshore environment (Saye & Pye, 2006). Sediment availability is the susceptibility of surface grains to entrainment by the wind (Kocurek & Lancaster, 1999); stabilizing factors such as early intergranular cements (for example, gypsum), vegetation cover, coarse-grained lags and elevated water tables all limit availability. Transport capacity is a measure of the potential sediment-carrying capacity of the wind. Together these factors define the sediment system state (Kocurek & Lancaster, 1999) which can be used as a predictor of when and where episodes of aeolian construction will occur.

Following Kocurek & Havholm (1993), three principal types of aeolian systems (Fig. 2) are recognized: (i) dry aeolian systems in which the water table and its capillary fringe are sufficiently far below the depositional surface that they have no effect on dune migration, sediment transport and deposition; (ii) wet aeolian systems in which the water table and its capillary fringe are at or near the depositional surface, so that changes in moisture play an important role in the style and pattern of sediment accumulation (Kocurek & Havholm, 1993; Mountney, 2012), and in which interdune areas are damp or wet (flooded) and characterized by clastic, biogenic and/or chemical sediments that are indicative of a near-surface water table; and (iii) stabilized aeolian systems in which factors such as vegetation, pedogenesis, permafrost or surface or near-surface cementation either episodically or continually act to stabilize the substrate while the system remains active overall, thereby encouraging aeolian construction and accumulation.




Fig. 2. Characteristics of dry, wet and stabilizing aeolian systems illustrating the role of aerodynamic configuration, water-table level and stabilizing agent as controls on accumulation space. After Kocurek (1998).

Aeolian accumulation to generate a body of strata requires a positive net sediment budget for which upstream sediment influx exceeds downstream outflux (Fig. 3). Special cases include aeolian accumulation in front of steep cliffs (e.g. Clemmensen *et al.*, 1997; Andreucci *et al.*, 2010a). By contrast, neutral budgets and negative budgets result in bypass and deflation (erosion), respectively. The positive net sediment budget required for aeolian accumulation needs either a downstream spatial decrease in the transport rate in response to airflow deceleration or a temporal decrease in flow concentration in response to a reduction in dune size over time (Rubin & Hunter, 1982; Kocurek & Havholm, 1993). One commonly recognized mechanism for the accumulation of migrating dunes and draas (mega-bedforms) is via bedform climbing, whereby the angle of climb (which for large bedforms might typically be only a few tenths of a degree) is determined by the ratio between the rate of downwind bedform migration and the rate of rise of the accumulation surface (Fig. 3). Climb at low angles means that only the basal parts of large bedforms typically accumulate to generate cross-stratified sets (Fig. 3). Nevertheless, accumulated, vertically stacked, cross-stratified sets recording the passage of multiple large bedforms are commonly each in excess of 10 m in thickness and some can attain thicknesses of >30 m (e.g. Mountney & Howell, 2000). The accumulation of sets via climbing and their composition of only the basal-most parts of the original bedforms from which they were constructed mean that ancient aeolian accumulations are biased representations of original aeolian systems because they are composed of assemblages of lithofacies arranged into architectural elements that typically record only those processes that operated on the lowermost flanks of the original bedforms; such processes typically differ from those that operated on the higher parts of bedforms (e.g. Eastwood *et al.*, 2012).

Interdune migration bounding surfaces separate packages of strata that represent the accumulated deposits of successive migrating aeolian dunes and adjoining interdunes; superimposition bounding surfaces record the style of juxtaposition of smaller dunes on larger draas, and the style of migration of the smaller forms over the larger forms; reactivation surfaces record episodic changes in dune or draa lee-slope configuration, including temporal changes in steepness or orientation (Rubin, 1987; Rubin & Carter, 2006). These bounding-surface types define and delineate architectural elements comprising packages of




Fig. 3. Spectrum of preserved dune and interdune architectures resulting from temporally and spatially invariable (i.e. static) aeolian system behaviour. The angle of bedform climb defines fields of accumulation and deflation, with bypass occurring when the angle of climb is zero. Within the field of accumulation, preserved sedimentary architecture is partly determined by the proportion of the accumulation surface covered by dunes. Accumulating dry aeolian systems typically require 100% dune cover whereby dunes have been constructed to a size where interdunes are reduced to isolated depressions between bedforms. Accumulating wet or stabilizing systems have less than 100% dune cover. The angle of climb is determined by the ratio of the vertical accumulation rate and bedform migration rate. The stratal configurations are scale-independent and can potentially occur in systems of any size; after Mounney (2012). Bedform spacing is the crest to crest (or toe to toe) distance between adjacent bedforms in an orientation perpendicular to the trend of elongate bedform crestlines; dune wavelength records the extent of a bedform in an orientation perpendicular to the trend of the bedform crestline and this may vary from a maximum dune wavelength to a minimum dune wavelength within one dune segment as a function of bedform sinuosity (Al-Masrahy & Mounney, 2013). Bedform spacing and dune wavelength will be the same for straight-crested dunes that lack intervening interdune flats.

aeolian dune and interdune strata that are themselves composed internally of various arrangements of lithofacies (Brookfield, 1977; Kocurek, 1991; Chrintz & Clemmensen, 1993; Fryberger, 1993). The geometry and arrangement of these architectural elements are determined by: (i) the scale and morphology of the original dunes and interdunes; (ii) the style of migration of the dunes and interdunes over both time and space; and (iii) the style of accumulation, which in many systems is controlled by the angle of climb (e.g. Mounney & Thompson, 2002), although in other systems it is known to be controlled by other mechanisms, including the infilling of local accommodation space between older remnant dunes (e.g. Langford *et al.*, 2008).

When and where the net sediment budget switches from positive to neutral or negative, aeolian accumulation ceases and bypass and defla-

tion commence, respectively. Both bypass and deflation result in the generation of supersurfaces (Kocurek, 1988) that cap underlying accumulations. Such accumulations define aeolian sequences and their bounding supersurfaces can be considered sequence boundaries (Fig. 4). Deflation operates either until the net sediment flux becomes neutral or positive again, or until it progresses down to the water table (Stokes, 1968), which limits further deflation. Supersurfaces of allogenic origin tend to be regional in extent and truncate other bounding-surface types of auto-genic origin, which themselves arise as a consequence of interdune migration and climb, bedform superimpositioning or bedform reactivation (Brookfield, 1977, Rubin, 1987; Mounney, 2006a). Some supersurfaces have been correlated laterally into adjoining non-aeolian environments where they merge into, for example, transgressive




Fig. 4. Spectrum of interdune geometries generated by variations in the frequency and magnitude of water-table change, the rate of dune migration and the net aeolian sediment budget. (A) Entrada Sandstone, Kocurek (1981a). (B) Navajo Sandstone (Herries, 1993). (C) and (D) Helsby Sandstone Formation (Mountney & Thompson, 2002). (E) Cedar Mesa Sandstone (Langford & Chan, 1988, 1989; Mountney & Jagger, 2004). (F) White Sands (Simpson & Loope, 1985; Loope & Simpson, 1992). Modified after Mountney & Thompson (2002).

marine units (Havholm *et al.*, 1993; Blakey, 1996; Blakey *et al.*, 1996; Rodríguez-López *et al.*, 2013). Many supersurfaces that bound episodes of aeolian accumulation are paraconformities (diastems) considered to represent long-lived hiatuses in accumulation: supersurfaces with associated sedimentary features such as large and closely spaced tree-size rhizoliths may take 10^4 to 10^5 years to form (Loope, 1985). Several authors have proposed that aeolian supersurface generation may occur as a result of Milankovitch-style orbital forcing operating with periodicities of 18 to 400 kyr (Loope, 1985; Clemmensen *et al.*, 1994; Mountney, 2006b; Jordan & Mountney, 2010, 2012; Rodríguez-López *et al.*, 2012a). For many systems, the amount of time represented by aeolian accumulations probably is significantly less than that represented by intervening supersurfaces (e.g. Loope, 1985); thus, many preserved aeolian successions probably represent only a fraction of the geological time over which the aeolian systems were active, and the preserved record is therefore highly fragmentary and potentially biased towards a specific set of formative processes.

Long-term preservation of aeolian accumulations in the ancient record requires that the body of strata is placed below some regional baseline, beneath which erosion does not occur (Kocurek & Havholm, 1993). Thus, the rate of generation of accommodation space and the rate at which aeolian accumulations fill that space is a fundamental control on preserved architectural style (e.g. Howell & Mountney, 1997).

Approaches to the theoretical modelling of aeolian dune and interdune successions commenced with the development of purely qualitative depositional models for aeolian systems. Many such models were devised in the 1970s and commonly recognised packages of aeolian dune and interdune lithofacies occurring as elements delineated by bounding surfaces (e.g. Brookfield, 1977). These models, which typically accounted for stratigraphic complexity in two spatial dimensions, are the so-called *static* aeolian depositional (or facies) models (Mountney, 2006a). One forward stratigraphic modelling approach to account for both spatial and temporal changes in aeolian architecture has led to the establishment of a conceptual framework for the classification of aeolian systems and their accumulated successions (Mountney, 2012). This framework identifies simple, *static* system architectures, that are generated by spatially and temporally invariable controls, but additionally identifies and models *dynamic* system architectures in which spatial and tempo-

ral changes in dune morphology, scale and style of migration and accumulation (for example, angle of climb) give rise to more complex preserved architectures.

EVOLUTION OF AEOLIAN SAND SYSTEMS THROUGH EARTH HISTORY

Archean and Proterozoic aeolian sand systems (Table S1)

The oldest known aeolian system is the 3.2 to 3.0 Ga Lower Moodies Group of the Swaziland Supergroup in South Africa (Table S1), which accumulated in a series of intramontane extensional basins through which simple barchan dunes migrated (Simpson *et al.*, 2012). The palaeogeographical distribution of Archean and Proterozoic aeolian systems was dictated by the worldwide geographical distribution of Archean and Proterozoic cratons and, due to their extreme age, the preserved global record of aeolian systems from these Eons is highly fragmentary. Palaeoproterozoic aeolian systems accumulated in intracratonic sag basins, intracratonic and intercontinental extensional rift basins (both during rift and thermal phases of basin evolution) and transtensional basins (e.g. Rainbird *et al.*, 2003; Master *et al.* 2010). Mesoproterozoic aeolian systems accumulated in intracratonic and intramontane basins, rift basins and transpressive strike-slip basins (e.g. Clemmensen, 1988; Martin & Thorne, 2002). Neoproterozoic aeolian systems accumulated in intracratonic rift basins (e.g. Grey *et al.* 2005; Sarkar *et al.*, 2011). From this, it is clear that the majority of preserved Precambrian aeolian systems are syn-rift depositional systems in which preservation of aeolian deposits occurred during the rifting phases of supercontinents because of the associated increase in accommodation space (Eriksson & Simpson, 1998).

Pre-vegetation Archean and Proterozoic periods were not subject to palaeoenvironmental conditions that were especially well-suited to aeolian sediment accumulation and preservation (Eriksson & Simpson, 1998). The absence of vegetation as a stabilizing agent favoured aqueous-reworking of pre-existing aeolian deposits, leading to their partial or total destruction, reworking and incorporation into a variety of coeval sedimentary environments (e.g. Tirsgaard & Øxnevad, 1998).

Proterozoic aeolian systems developed in association with a variety of coeval depositional systems and led to fluvial–aeolian interactions, such

as those from the Palaeoproterozoic Makgabei Formation (Waterberg Supergroup, South Africa, Eriksson *et al.*, 2000; Simpson *et al.*, 2002), Amarakoo Formation (Wharton Group, Canada, Rainbird & Hadlari, 2000; Rainbird *et al.*, 2003) and Thelon Formation (Barrenland Group, Canada, Rainbird *et al.*, 2003). Marine reworking of Precambrian aeolian deposits was apparently a widely occurring process along Proterozoic coastlines; documented examples include the Palaeoproterozoic Quartzite Member, Mcheka Formation, Zimbabwe (Master *et al.*, 2010), the Whitworth Formation, Haslingden Group, Australia (Simpson & Eriksson, 1993) and the Neoproterozoic Venkatpur Sandstone, India (Chakraborty, 1991).

Comparison of Proterozoic and Phanerozoic erg systems reveals a general trend towards the preservation of more complex aeolian systems during the Proterozoic. The general atmospheric circulation pattern influenced by palaeogeographical changes, palaeoland mass distributions and associated orogenic buildups, and the particular properties and characteristics of Archaean and Proterozoic atmospheres could have had a different effect on aeolian transport compared to equivalent processes that operated during the Phanerozoic. Studies by Han *et al.* (2014) have demonstrated that, for a particular wind speed, the ability of the air flow to transport sand decreases with lower air density; however, under the same conditions, the saltation height increases. Taking into account that recent works like that of Som *et al.* (2012) have concluded that the density of the 2.7 Ga atmosphere was less than twice modern levels, it is possible that changes in air density over geological timescales could have influenced aeolian transport mechanisms, and this might be recorded by the predominance of different aeolian bedforms at different times. For example, documented examples of co-sets of strata interpreted to represent the preserved accumulations of draa-scale bedforms are numerous for Precambrian successions. Furthermore, the preservation of very coarse-grained (siliciclastic) Precambrian aeolian successions (for


example, the Egalapemta Member, Mesoproterozoic, India; Biswas, 2005) is noteworthy. The specific dynamic configuration of the Precambrian atmosphere and its interaction with sediment grains could explain the occurrence of simple but giant transverse dunes with maximum preserved set thicknesses (more than 50 m thick), such as the single aeolian dune cross-bedded set recorded from the Late Neoproterozoic McFadden Formation (Western Australia; Grey *et al.*, 2005).

The absence of Phanerozoic cold (periglacial) aeolian dunefields compared with their occurrence in Precambrian times is notable. The Neoproterozoic Bakoye 3 Formation, Bakoye Group from Mali (Deynoux *et al.*, 1989) and the Neoproterozoic Whyalla Sandstone from Australia (Williams, 1998) constitute the only two examples of Pre-Cenozoic periglacial dunefields. This scarcity of periglacial dunefields in the fossil record could be a result of misinterpretation of the particular palaeoclimate setting in which some Precambrian and Phanerozoic aeolian systems formed. A re-evaluation of this topic is needed. It is known that glacial latitudes have changed through time (see Evans, 2003). The possibility that cold deserts could have formed in association with glacier fronts even during the Phanerozoic should be considered. Particular attention should be paid to the latitudinal variation of the equilibrium-line altitude (ELA) as a control of glaciation through time (e.g. Isbell *et al.*, 2012).

Palaeozoic aeolian sand systems (Table S2)

During the Cambrian, ergs were located in the main land masses in the Southern Hemisphere (Fig. 5A). The Backbone Ranges Formation (Mackenzie Mountains, Canada; MacNaughton *et al.*, 1997) and the Wonewoc Formation (Wisconsin and Minnesota, USA; Dott *et al.*, 1986; Runkel *et al.*, 1998) accumulated in southern Laurentia where interaction between ergs and coastal systems adjacent to the Iapetus Ocean occurred (Fig. 5A). The Amin Formation and the Lower Haima Group in Oman (Milson *et al.*, 1996) and the Lower Roan Formation in Zambia

Fig. 5. Palaeogeographic distribution of siliciclastic aeolian sand systems during: (A) Late Cambrian. During the Cambrian, ergs were located in main land masses at the Southern Hemisphere (Laurentia and north-eastern Gondwana), under the influence of easterly trade winds between the Equator and 30° palaeolatitudes. (B) Middle Ordovician. Main ergs of Cambro-Ordovician and Ordovician age were located in Laurentia and eastern Gondwana under the activity of trade winds of subtropical low pressure systems. (C) Middle Silurian. Main Silurian and Siluro-Devonian aeolian deposits are located in western/central Australia, located at the southern palaeo-hemisphere desert zone. (D) Early Devonian. Palaeogeographic maps from Scotese (2001) PALEOMAP Project. For legend see Figure 6D.



(Annels, 1989) accumulated as inland sandy deserts in Gondwana (Fig. 5A). Many Cambrian ergs accumulated under the influence of easterly trade winds, between the Equator and 30° palaeolatitude (Fig. 5A; see Dott *et al.*, 1986).

Cambro-Ordovician and Ordovician ergs were less widespread than those of Cambrian age (Fig. 5B). Such aeolian systems again developed under the influence of active trade winds of subtropical high-pressure systems in land masses of the southern palaeo-hemisphere in Laurentia and Gondwana (Fig. 5B) (for example, the Pedra Pintada Formation/Alloformation; Paim & Scherer, 2007; de Almeida *et al.*, 2009). In particular, aeolian dunefields developed in southern Laurentia record marine–aeolian interactions characterized by complex associations of facies of both aeolian and aqueous origin, as is the case for the Cambro-Ordovician Nepean Formation (Postdam Group, Canada and USA; Malhame, 2007) and the Ordovician St. Peter Sandstone (Minnesota and Wisconsin, USA; Dott *et al.*, 1986). In particular, the St. Peter erg succession records palaeowinds that are in agreement with more general reconstructions of the southern palaeo-trade wind belt (Dott *et al.*, 1986).

Silurian and Siluro-Devonian aeolian successions are few in number. The main preserved systems accumulated in Western (Perth-Carnarvon Basin) (Trewin & Fallick, 2000) and Central (Amadeus Basins) Australia (Fig. 5C) (Shaw *et al.*, 1991). The Swanshaw Sandstone Formation of Scotland constitutes a mixed aeolian–fluvial succession developed in the transtensional Lanark Basin (Smith *et al.*, 2006). These Silurian and Siluro-Devonian systems developed in the southern palaeo-hemisphere desert belt.


During the Devonian, the assembly of Laurussia in response to the final stages of the plate collisions of the Caledonian Orogeny, and the northward migration of Gondwana, led to an increase of land masses present at subtropical latitudes that were subject to the influence of the southern palaeo-hemisphere desert belt (Fig. 5D). This palaeogeographical configuration enabled the construction, accumulation and preservation of several major aeolian dunefield systems (Fig. 5D).

The most representative Devonian aeolian systems are those forming part of the Old Red Sandstone of North West Europe (e.g. Browne *et al.*, 2002; Morrissey *et al.*, 2012) which accumulated in extensional basins formed as a result of the collapse of the overthickened crustal belt resulting from Caledonian compressional tectonics (McClay *et al.*, 1986). The Old Red Sandstone exhibits a variety of aeolian facies, many of which record wind–water interaction processes (for example, the Middle Devonian Yesnaby Sandstone Group, Lower Old Red Sandstone Supergroup, Scotland; Trewin & Thirlwall, 2002).

Devonian ergs are characterized by a variety of aeolian facies including aeolian sandsheet successions (for example, the Lower Clair Group, Clair Basin, UK; Nichols, 2005), aeolian dunefield successions composed of transverse dune deposits (for example, the Slieve Mish Group, Ireland; Horne, 1971), barchanoid dune deposits (for example, the Devonian of Scotland; Allen & Marshall, 1981) and draa deposits (for example, the Kilmurry Sandstone Formation, Ireland, Dodd, 1986; the Eday Sandstone, Eday Group, Scotland, Marshall *et al.*, 1996). Devonian aeolian dunefield successions with draa and barchanoid dune deposits are preserved in north-east Greenland and these demonstrate aeolian interaction with ephemeral streams and terminal fans (see Olsen & Larsen, 1993). Other Devonian aeolian systems have been recorded from Antarctica (New Mountain Sandstone; Gilmer, 2008) and Australia (for example, the Langra Formation, Jones, 1972; the Tandalgoo Sandstone, Thornton, 1990).

The majority of Carboniferous aeolian systems are Pennsylvanian in age, with several spanning the Pennsylvanian–Permian boundary. Some examples of early Carboniferous aeolian systems include the Devonian to Mid-Carboniferous Khusayyayn Formation in Saudi Arabia (Stump & Van der Eem, 1995), the recently recognized aeolian systems of the Late Mississippian Loyalhanna Member, Pennsylvania, USA (the Mauch Chunk Formation and Appalachian Formation; Swezey *et al.*, 2012) and the Devonian–Mississippian Harder Bjerg Formation in Greenland (Olsen & Larsen, 1993; Fig. 6A).

Fig. 6. Palaeogeographic distribution of siliciclastic aeolian sand systems during: (A) Early Carboniferous. Early Carboniferous aeolian systems were located in southern Laurussia and at the northern margin of Gondwana. (B) Late Carboniferous. Late Carboniferous–Permian aeolian systems were located in Pangea supercontinent both at Northern and Southern Hemispheres. Other aeolian systems developed in northern Gondwana (Bolivia and Argentina). (C) Late Permian. During Permian times extensive erg systems developed on Pangea, favoured by the location of this supercontinent in subtropical latitudes under the influence of the southern and northern palaeo-hemisphere desert zones. Palaeogeographic maps from Scotese (2001) PALEOMAP Project. (D) Legend is for Figures 5–8.



During the Variscan Orogeny, Gondwana and Laurussia collided creating the Supercontinent Pangaea; Pennsylvanian to Permian aeolian systems were constructed in both the Northern and Southern Hemispheres (Fig. 6B). The main systems crop out in North and South America, with well-developed examples including the Early Pennsylvanian Juruá Sandstone Formation (the Solimões Basin, Brazil; Elias *et al.*, 2007), the Pennsylvanian Tyrwhitt and Tobermory Sandstone Formations (the Rocky Mountain Supergroup, Canada; Stewart & Walker, 1980) and the Pennsylvanian–Middle Permian Cangapi Formation (the Cuevo Group, Tarija Basin, Bolivia and Argentina; Hernández & Echevarría, 2009), the Late Carboniferous–Early Permian Patquía Formation (the Paganzo Group, Paganzo Basin, Argentina; Caselli & Limarino, 2002; Geuna *et al.*, 2010). In Saudi Arabia, the Carboniferous–Permian Unayzah Formation constitutes an economically important gas reservoir succession (Melvin & Heine, 2004; Melvin *et al.*, 2010).

The best-known Pennsylvanian–Permian ergs are reported from the USA. The ‘Tensleep Complex’ (the Tensleep Sandstone, Casper Formation, Quadrant Sandstone; e.g. Peterson, 1988), the Honaker Trail Formation (e.g. Williams, 2009) and the Cutler Group, including the lower Cutler beds (e.g. Jordan & Mountney, 2010, 2012; Wakefield & Mountney, 2013), the Rico Formation (e.g. Loope, 1985; Chan & Kocurek, 1988) and the Weber Sandstone (e.g. Doe & Dott, 1980; Driese, 1985) are all successions that exhibit well-exposed examples of central-erg and erg-margin systems (see Blakey *et al.*, 1988, for further examples).

During the Permian, the construction of extensive erg systems across large parts of Pangaea was favoured by the location of land masses of this supercontinent in subtropical latitudes under the influence of the southern and northern palaeo-hemisphere desert zones (Fig. 6C). The best-known Permian aeolian systems are the Rotliegend Group (Rotliegendes) of the North Sea and North-west Europe, and the Permian aeolian systems from the USA (Fig. 6C). Major reserves of gas (and some oil) exist in Permian Rotliegend desert sandstone hydrocarbon reservoirs of North-west Europe, in particular in the Southern Permian Basin of the North Sea and some localities in the Northern Permian Basin (Glennie, 1970, 1972, 1998; Glennie & Buller, 1983).

In the Southern Permian Basin of the North Sea, aeolian dune deposits accumulated between

wadi channels originating from the Variscan Highlands and the extensive sabkha and desert lake located southwards of the Ringkøbing-Fyn High (Glennie, 1972). Reconstructed aeolian dune types of the Rotliegend Group include transverse-crescentic dunes, barchans, longitudinal/linear and star dunes (for example, the Brodick Beds; Clemmensen & Abrahamsen, 1983; the Leman Sandstone Formation, Sweet, 1999; the Penrith Sandstone, Turner *et al.*, 1995; Lovell *et al.*, 2006), as well as interdraa, draa-plinth and draa-centre deposits (for example, Yellow Sands; Clemmensen, 1989; Chrintz & Clemmensen, 1993). Complex wind patterns resulted in the construction of barchanoid draa with superimposed oblique crescentic and linear dunes (for example, the Bridgnorth Sandstone Formation, UK; Steele, 1981; Benton *et al.*, 2002).

Permian sedimentary basins of the USA contain extensive and complex aeolian depositional systems and record a variety of facies and processes (see Blakey *et al.*, 1988, for compilation). Examples of these Permian aeolian units include the Schnebly Sandstone Formation, which comprises deposits of an aeolian dunefield associated with evaporite and carbonate deposits (Blakey & Middleton, 1983; Blakey, 1990), and the Lyons Sandstone Formation with deposits of parabolic dunes and blowout-type interdunes (McKee, 1979).

Several Permian units in the USA preserve complete examples of central-erg sequences that demonstrate evidence for a complex merging relation with marine erg-margin systems. Examples include the White Rim Sandstone (e.g. Chan, 1989; Tewes & Loope, 1992; Kamola & Huntoon, 1994), the De Chelly Sandstone (e.g. Blakey, 1990; Stanesco, 1991), the Yeso Formation (Mack & Dinterman, 2002) and the Upper Minnelusa Formation (e.g. Fryberger, 1984, 1993).

The Permian Coconino Sandstone constitutes the accumulation of an inland dry erg system formed by climbing barchans or barchanoid-ridge and transverse dunes (e.g. Blakey & Middleton, 1983; Blakey, 1990, 1996). Some Permian ergs display examples of fluvial systems reworking the aeolian sands; this is the case for the Rush Springs Sandstone (the Whitehorse Group; Kocurek & Kirkland, 1998; Poland & Simms, 2012), the Cedar Mesa Sandstone of the Paradox foreland basin (Langford & Chan, 1988, 1989; Mountney & Jagger, 2004; Mountney, 2006a; Langford *et al.*, 2008) and the overlying Organ Rock Formation (Cain & Mountney, 2009, 2011).

Permian aeolian systems developed in the Southern Hemisphere in Pangaea (Fig. 6C) include the Pirambóia Formation in Brazil (Paraná Basin; Dias & Scherer, 2008), the Buena Vista Formation in Uruguay (Northern Uruguayan Basin; Goso *et al.*, 2001) and the Permian aeolian systems of Argentina from the retroarc Paganzo Basin (for example, the Andapaico Formation and the De la Cuesta Formation; Spalletti *et al.*, 2010; Correa *et al.*, 2012).

Mesozoic aeolian sand systems (Table S3)


Throughout much of the Triassic, widespread aeolian desert and semi-desert conditions prevailed across much of northern Pangaea. The majority of Triassic ergs were located in equatorial to mid-latitudes in the Northern Hemisphere and most of these appear to be aligned following a north–south trend close to the eastern margin of Northern Gondwana (Fig. 7A). Triassic erg systems of north-eastern Pangaea include the Buntsandstein of Europe, which is characterized by a thick accumulation of red beds that record a variety of aeolian and mixed aeolian–fluvial–lacustrine successions (e.g. Clemmensen, 1985; Mader, 1985; Mader & Laming, 1985; Tietze *et al.*, 1997). The Triassic Buntsandstein facies in the north-eastern Iberian Chain (central eastern Spain), previously considered to be fluvial in origin, is now known to contain an evolving erg system (Soria *et al.*, 2011) which comprises a succession that records the transition from a wadi belt, via an inner erg-margin, to a central-erg system.

The equivalent lithostratigraphic unit to the Buntsandstein in the UK and Ireland is the Sherwood Sandstone Group ('New Red Sandstone') which is present in a series of rift basins in both onshore and offshore settings (Fig. 7A; Brookfield, 2004, 2008; Tyrrell *et al.*, 2009). Aeolian dunefields were mostly characterized by bedforms of modest size, many with damp or wet interdunes controlled by the water table, as recorded, for example, by the Wilmslow Formation (Øxnevad, 1991; Bloomfield *et al.*, 2006) and the Helsby Formation (Mountney & Thompson, 2002; Bloomfield *et al.*, 2006) of the Cheshire Basin. In Scotland, the Hopeman Sandstone probably straddles the Permian–Triassic boundary and is characterized by deposits of the preserved remnants of a series of star dune and draa bedforms representing a small fragment of what is inferred to have been a very extensive dry aeolian system (Clemmensen, 1987; Glennie

& Hurst, 2007; Hurst & Glennie, 2008). In the subsurface of the East Irish Sea Basin, Triassic aeolian deposits form important reservoirs for gas (Cowan & Boycott-Brown, 2003; Meadows, 2006).

Triassic ergs constructed close to the palaeo-equator (for example, the Oukaimeden Sandstone Formation, Morocco) record small aeolian dunes developed on floodplains of ephemeral fluvial systems (Fabuel-Perez *et al.*, 2009; Mader & Redfern, 2011). Other Triassic ergs are located close to the western margin of Northern Pangaea and examples include the Nugget Sandstone of Utah and Wyoming (Fig. 7A) (Sprinkel *et al.*, 2011) which may be, at least in part, of Lower Jurassic age.

Thick and geographically widespread Jurassic aeolian desert erg successions of the Colorado Plateau region are extensively documented and are arguably the most intensely studied of all aeolian successions. Many authors have considered these successions collectively in terms of the regional palaeogeographical, palaeoclimatic and palaeotectonic setting (Fig. 7B and C; e.g. Kocurek & Dott, 1983; Blakey *et al.*, 1988; Dickinson & Gehrels, 2003; Loope *et al.*, 2004). The best-known Jurassic aeolian successions of the south-western United States include the Wingate and Navajo sandstones (and stratigraphic equivalents) of the Lower Jurassic Glen Canyon Group, and the Page and Entrada sandstones (and equivalents) of the Middle Jurassic San Rafael Group. The Wingate Sandstone represents a largely dry aeolian system representative of an erg-centre setting (with compound-draa development) and also demonstrates styles of interaction with fluvial deposits of the Moenave Formation in its erg-margin setting (Clemmensen & Blakey, 1989; Clemmensen *et al.*, 1989; Tanner & Lucas, 2007). The Navajo Sandstone of the Glen Canyon Group is one of the most intensely studied sedimentary formations of any type and is well-exposed across much of the Colorado Plateau region, where it attains a thickness of nearly 700 m in south-western Utah. The succession represents the preserved remnant of a giant erg that was present across much of the western part of Pangaea (Fig. 7B; Hunter & Rubin, 1983; Chan & Archer, 2000); this system was subject to seasonal wind reversals associated with annual monsoons that occurred each summer when more humid and cooler conditions prevailed and wind reversal occurred (Loope & Rowe, 2003; Loope *et al.*, 2008). The Page Sandstone of Arizona and southern Utah repre-



sents accumulation in an erg system close to the margin of an interior seaway; the system is composed of vertically stacked, progradational erg sequences that overlie marine deposits of the Carmel Formation such that the two units inter-tongue (Jones and Blakey, 1993; Havholm & Kocurek, 1994; Dickinson *et al.*, 2010). The Entrada Sandstone of the San Rafael Group is exposed extensively across much of the Colorado Plateau region and represents the accumulated deposits of a coastal to inland aeolian system that was characterized by a complex arrangement of aeolian dune, damp and wet interdune, and sabkha elements (Kocurek, 1980, 1981a,b; Crabaugh & Kocurek, 1993; Crabaugh & Kocurek, 1998). Relic dune topography is preserved in places at the top of the succession as a result of later marine transgression (Benan & Kocurek, 2000).

The Upper Jurassic Norphlet Sandstone represents the accumulated deposits of a major aeolian erg succession that is known principally from the subsurface of Alabama, the shallow-water Gulf of Mexico around Mobile Bay, and further offshore in the deep-water part of the Gulf of Mexico, where it forms a major oil reservoir (Taylor *et al.*, 2004; Mankiewicz *et al.*, 2009; Ajdukiewicz *et al.*, 2010). Numerous aeolian successions of Jurassic age are documented from South America and especially from Brazil. Examples include the Pedreira Sandstone of the Paraná Basin in Brazil, which is characterized by climbing aeolian dune sets with intervening damp and wet interdune units (Nowatzki & Kern, 2000) and the Guar Formation of south-

ern Brazil, which records composite crescentic aeolian dune sets and cosets, and aeolian sand-sheet elements interbedded with distal flood deposits and fluvial channel-fill elements (Scherer & Lavina, 2006).

The backarc Neuqun basin of Argentina records a series of Jurassic aeolian successions. The Lotena Formation preserves a record of aeolian–fluvial interactions (Veiga *et al.*, 2011a). A tectonic inversion during the Late Jurassic led to the desiccation of the entire basin, giving rise to a complex array of continental facies for which aeolian deposits form a major part (Spalletti & Veiga, 2007, Spalletti *et al.*, 2011). In the southern part of the basin, an upward vertical transition from fluvial-dominated to aeolian-dominated deposition, probably arising from a climatic shift to drier conditions, is recorded as part of the Kimmeridgian Quebrada del Sapo and Tordillo formations (Zavala *et al.*, 2005a; Veiga & Spalletti, 2007). The Tordillo Formation represents migration and accumulation of transverse and barchan dunes in a style that generated a complex hierarchy of internal bounding surfaces within a largely dry aeolian system in which only thin dry interdune elements accumulated (Zavala *et al.*, 2005a). To the east, aeolian accumulation was more significant and led to the preservation of a *ca* 300 m thick sequence of mainly aeolian deposits of the Sierras Blancas and Catriel formations, which include deposits of dune, wet and dry interdune and aeolian sand-sheet elements (Maretto *et al.*, 2002; Spalletti *et al.*, 2011). The Piramboia Formation of Entre Rios Province, Argentina, is an aeolian

Fig. 7. Palaeogeographic distribution of siliciclastic aeolian sand systems during: (A) Early Triassic. Triassic erg systems occupied a broad belt across the tropical latitudes of northern Pangea in areas now occupied by Europe and North America. Other Southern Hemisphere erg systems are recorded from Gondwana (S. Africa and India). (B) Early Jurassic. The main erg systems of the Early Jurassic include the geographically widespread accumulations of the Glen Canyon Group in the western USA (including the Wingate, Navajo and Page sandstones) and accumulations in the Paran and Neuqun basins. (C) Late Jurassic. The main erg systems of the Late Jurassic include the geographically widespread accumulations of the San Rafael Group in the western USA (including the Entrada Sandstone) and accumulations in the Paran Basin (Brazil) and southern Africa. (D) Late Cretaceous. Cretaceous erg systems were geographically widely distributed in tropical and mid-latitudes in both the Northern and Southern Hemispheres, with major dunefields occupying SW Africa and Central Brazil prior to and immediately following the onset of break up of western Gondwana. Other major ergs were located in northern China and Spain. Palaeogeographic maps from Scotese (2001) PALEOMAP Project. For each map, the aeolian successions shown are not all necessarily of the same age and, therefore, were not necessarily all active at the same time. Furthermore, the global palaeogeographies depicted in each map might not necessarily be entirely accurate for aeolian successions that are slightly older or younger than the age shown in the maps. For example, the early Cretaceous Botucatu Sandstone of the Paran Basin of Brazil and the Etjo Sandstone of NW Namibia are considered to represent preserved portions of the same erg system that developed prior to the onset of opening of the South Atlantic, yet the palaeogeographic map shown depicts an interval in the Late Cretaceous, shortly after the onset of the opening of the south Atlantic. For legend see Figure 6D.

unit considered to be primarily of Lower Jurassic age (Silva & Fernandez, 2004).

The Lower Jurassic Clarens Formation, which forms a unit of the Karoo Supergroup in South Africa, records a progressive upward transition from the deposits of a wet aeolian system that developed alongside coeval ephemeral fluvial systems to a dry aeolian system dominated by stacked cross-bedded aeolian dune sets (Bordy & Catuneanu, 2002; Holzförster, 2007).

Cretaceous aeolian successions, together with those that probably span the Jurassic–Cretaceous boundary, are numerous in South America and many have been the focus of detailed study over several decades (Fig. 7D). The Botucatu Formation of the Paraná Basin (São Paulo and Paraná states, Brazil) – which was originally thought to be Triassic in age (Bigarella, 1979) – has lateral equivalents in Parnaíba Basin of northern Brazil and is a near-equivalent of the Bauru, Guará, Sambaíba, Sanga do Cabral and Piramboia formations, as well as of the Etjo Sandstone in Namibia (Mountney *et al.*, 1999a,b), and possibly the Kudu Formation, offshore Namibia (Mello *et al.*, 2011). Relic aeolian dune forms and degraded topography are preserved at the top of the succession where it is overlain by flood basalts of the Serra Geral Formation and other flood basalts related to the Etendeka-Paraná Large Igneous Province (Scherer, 2002; Waichel *et al.*, 2008). The Serra Geral Formation records the exceptional preservation of relic aeolian dune topography of a dry aeolian system by flood basalts including various types of completely preserved dunes and sand-deformation features, including sand diapirs and peperite-like breccia.

In the backarc Neuquén Basin of west-central Argentina, Lower Cretaceous sandy aeolian accumulations are numerous and mainly related to lowstand periods and to the possible disconnection of the basin from the proto-Pacific Ocean (Howell *et al.*, 2005). These successions constitute important conventional oil and gas reservoirs. Aeolian deposits have been described from part of the proximal system of the Valanginian Mulichinco Formation in the subsurface of the basin (Zavala *et al.*, 2005b) and, more marginally, as part of environments of fluvial–aeolian interaction (Schwarz *et al.*, 2011). One of the best-described aeolian systems in the basin is the Hauterivian Avilé Member of the Agrío Formation (Rossi, 2001; Veiga *et al.*, 2011b). Within this non-marine unit, aeolian deposits are locally important and record a com-

plex vertical evolution related to high-frequency climatic changes and to the development of multiple supersurfaces associated with aeolian deflation and fluvial flooding (Veiga *et al.*, 2002). Finally, the Baramian Lower Troncoso Member of the Huitrín Formation is characterized by the transition from fluvial to aeolian deposits (Veiga *et al.*, 2005). For both the Avilé and Troncoso members, marine inundation of the dunefields following transgression led to the preservation of relic dune topography, as well as to the development of a complex set of facies related to the deformation and reworking of the aeolian sands during the transgression (Strömbäck *et al.*, 2005; Veiga *et al.*, 2011b). Aeolian deposits have also been described in the Upper Cretaceous record of the Neuquén Basin as part of the Neuquén Group (Sánchez *et al.*, 2008).

In Africa, the Lower Cretaceous Etjo Sandstone Formation of north-west Namibia is a predominantly dry aeolian system in which relic aeolian dune bedforms with up to 100 m of topographic relief have been preserved following inundation by flood basalts of the Etendeka Igneous Province (Mountney *et al.*, 1999a,b; Mountney & Howell, 2000; Howell & Mountney, 2001). The lower part of the succession records exceptionally thick examples of simple cross-bedded sets of aeolian dune origin (individual simple sets up to 52 m thick), with preservation probably having been enabled by the migration of a large dune into a pre-existing topographic depression. Aeolian sandstone occurs interleaved with flood basalts at multiple levels within the upper part of the succession, which forms the lower part of the overlying Etendeka Group Large Igneous Province (Jerram *et al.*, 1999a,b, 2000a,b).

Various formations composed of Cretaceous strata of aeolian dune origin are present in several basins of China, including many in the Gobi Desert region of Inner Mongolia. Cretaceous aeolian dune accumulations are recorded from the Sichuan, Ordos, Kuche, Tarim and the Kuqa basins, and especially in Inner Mongolia and surrounding regions. The preserved aeolian dunefield deposits preserve evidence for the development of both dry and wet (water-table controlled) aeolian systems (Xie *et al.*, 2005; Jiang *et al.*, 2008). The Upper Cretaceous (Campanian) Djadokhta Formation of the Ulan Nur Basin and the area around Tugrikiin Shiree and Ukhaa Tolgod (Nemegt Basin, Mongolia) represent dunefields that experienced heavy rainfall events resulting in the development of

perched water tables, early calcite cementation and dune collapse due to sediment gravity sliding (Jerzykiewicz *et al.*, 1993; Loope *et al.*, 1999; Seike *et al.*, 2010).

In Europe (Spain), the mid-Cretaceous Iberian Desert System, represented by the Utrillas Group (that includes the previously known middle and upper parts of the Escucha Formation and the whole Utrillas Formation) developed from the early Albian to the early Cenomanian along the western Tethyan margin (Iberian Basin, eastern margin of Iberia) between the Tethys Ocean (to the east) and the highland Variscan Iberian Massif (to the west) over an area of more than 20 000 km² (Fig. 7D; Rodríguez-López, 2008; Rodríguez-López *et al.*, 2008). The mid-Cretaceous Iberian Desert System displays a tripartite spatial configuration: a back-erg characterized by aeolian–fluvial (wadi) interactions, a central-erg characterized by thick accumulation of linear draa, other compound-draa sandstones and desert roses, and a fore-erg in which the interaction between compound aeolian dunes (draas) and coastal sedimentary environments (lagoons, tidal creeks, tidal deltas and marshes) occurred (Rodríguez-López *et al.*, 2006, 2008, 2010, 2012a). The sedimentary record of this desert basin displays different erg sequences bounded by supersurfaces (Rodríguez-López *et al.*, 2013).

Palaeogene aeolian sand systems (Table S4)

Palaeogene erg systems have been the subject of only relatively modest investigation, mainly as part of regional studies; it is therefore difficult to draw conclusions regarding their distribution and development. Only one example of aeolian accumulation has been described for the Palaeogene of Europe (Fig. 8A) and this corresponds to the Sables de Fontainebleau Formation (or Fontainebleau Formation) of early Oligocene age (Alimen, 1936). This unit is part of the fill of the Paris Basin and is composed of a 50 to 70 m thick succession of clean, fine-grained, well-sorted sand arranged in accumulations expressed as elongated ridges and is thought to represent the preserved topography of an ancient coastal barrier system (Thiry *et al.*, 1988; Cojan & Thiry, 1992).

Palaeogene aeolian systems of North America are restricted to those of the Oligocene of the western USA (Fig. 8A). The most important example corresponds to the ‘Chuska Erg’, an extensive sand sea (*ca* 140 000 km²) developed

in the uplifted Colorado Plateau between 33.5 Ma and 27 Ma (Lucas & Cather, 2003; Cather *et al.*, 2008). The accumulated record of this sand sea (known as the Nabora Pass Member of the Chuska Sandstone) attains a maximum thickness of 535 m and records the northerly migration of transverse dunes (Cather *et al.*, 2008). Also in the western USA, in the Great Plains of South Dakota, Nebraska and Wyoming, volcanoclastic aeolian deposits have been described as part of the Brule Formation of the White River Group (Tedford *et al.*, 2005).

Few aeolian successions have been described for the Palaeogene of Africa and these are mainly of Middle to Upper Eocene age (Fig. 8A). The Hadida Formation, which developed in the Tindouf-Ouarzazate Basin of Morocco, includes medium-grained, cross-bedded sandstone that occurs intercalated in a >300 m thick sequence mainly composed of gypsiferous mudstones (Swezey, 2009; Tesón *et al.*, 2010); this succession is regarded as the earliest record of the Saharan system (Swezey, 2006). Apart from these proto-Saharan deposits, sandy aeolian successions have been described as part of the Palaeocene fill of the Congo Basin in the West African margin of Gabon in the form of deposits originally described as the ‘Gres Polymorphes’ (De Ploey *et al.*, 1968) that comprise a 180 m thick succession of cross-stratified sandstones with individual sets several metres thick interpreted as aeolian deposits (Batéké Sands, Séranne *et al.*, 2008). Aeolian deposits have also been described in the Fayum District in Egypt as part of the Qasr El-Sagha Formation (El-Fawal *et al.*, 2011). These include a 45 m thick succession of Middle to Upper Eocene age, previously described as channelized delta plain deposits (Bown & Kraus, 1988), but more recently reinterpreted as part of a desertification phase that caps a prograding deltaic system (El-Fawal *et al.*, 2011).

The only example of Cenozoic aeolian accumulation from Oceania comes from southern Australia. The upper portion of the Middle to Upper Eocene Ooldea Sand (Barton Sand) represents a barrier dune complex developed during the transgression of the Eucla Basin (Hou *et al.*, 2006).

Neogene aeolian sand systems (Table S4)

Neogene sandy aeolian systems are relatively common and have been described worldwide (Fig. 8B and C). Most of these systems owe their




Fig. 8. Palaeogeographic distribution of siliciclastic aeolian sand systems during: (A) Eocene. Palaeogene aeolian systems are scarce. Larger accumulations are located in the Colorado Plateau in western USA. African systems may be the precursors of Quaternary deserts and coastal aeolian systems are described in France and Australia. (B) Miocene. Miocene ergs are more numerous and mostly located in western South America in the Andes foreland. Western North America systems are also important. In Africa and the Middle East ergs are related to similar conditions as in the Quaternary. (C) Last Glacial Maximum (18 ka). Pliocene ergs are related in Europe to the onset of Northern Hemisphere glaciation and the development of stronger westerlies. African systems indicate even drier conditions as in present times (for example, Mega Kalahari). Palaeogeographic maps from Scotese (2001) PALEOMAP Project. For legend see Figure 6D.

origin to local climatic and tectonic factors. However, as the position of most major continental landmasses has not changed significantly since the Miocene, many Neogene aeolian systems have apparently been controlled by climatic conditions similar to those experienced by present-day desert systems, and such successions therefore constitute the precursors of some of the most important Quaternary aeolian systems, as in the Sahara, Kalahari and Namib sand seas.

Only one example of a Miocene aeolian accumulation in Europe has been identified. It corresponds to the Vale de Chelas Sands in the Lower Tejo Basin of Western Iberia, a 10 m thick succession of cross-bedded sandstones of aeolian origin, interpreted as a coastal system (Telles Antunes *et al.*, 1999; Pais *et al.*, 2012). Pliocene aeolian deposits in Europe have been described in more detail and are mainly related to the onset of the Northern Hemisphere glaciation that led to stronger westerly winds (Fig. 8C). In central Spain, the Middle to Upper Pliocene Escorihuela Formation records the accumulation and preservation of an aeolian dunefield related to synsedimentary activity of normal faults in a syn-rift environment. This system is dominated by the interaction between constructive and destructive episodes related to high-frequency climatic changes (Rodríguez-López *et al.*, 2012b). Accumulation during the Middle to Upper Pliocene is also recorded in the northern Apennines of Italy (Fig. 8C), and is associated with extensional tectonics and the alternation between relatively more humid and more arid episodes in the Valdarno Basin (Ghinassi *et al.*, 2004). Here, deposits of the Rena Blanca Sand Unit are dominated by the superimposition of wetting–drying–wetting cycles that record high-frequency climatic oscillations, each apparently of *ca* 40 ka duration (Ghinassi *et al.*, 2004).

In the USA, several Miocene aeolian systems have been described and are mainly controlled by local tectonic conditions associated with warm and dry climatic conditions. In western and central USA, some local systems have developed related to extensional basins, such as the Zia Formation of the Santa Fe Group in the Albuquerque Basin (Galusha & Blick, 1971) and the Ojo Caliente Sandstone of the Tesuque Formation in the La Española Basin (Koning *et al.*, 2004), both related to the large structure of the Río Grande Rift. These systems give rise to locally thick successions (up to 160 m) related

to the development of dunefields that were strongly influenced by local conditions. Aeolian deposits have also been described in the High Plains of the USA, including the Early Miocene Arikaree Formation (Bart, 1977) in south-east Wyoming, where large-scale, cross-bedded sandstones have been related to the accumulation of barchan and transverse dunes. Finally, an aeolian origin has also been reported for a *ca* 100 m thick succession of the Comondú Group in Baja California, Mexico, related to the infill of the forearc basin developed between the Late Oligocene and Early Miocene (Umhoefer *et al.*, 2001).

Neogene aeolian accumulations in South America are related to the complex evolution of the Andes. The compressional regime in the western margin of South America led to the development of a complex foreland with multiple basins that formed important sites of accommodation that were themselves subject to an arid local climate regime. This resulted in the accumulation of several aeolian units, some with local names that record this stage of evolution, especially during the Miocene. For instance, the Petaca Formation in southern Bolivia (Uba *et al.*, 2005) and the Aguada Member of the Chacras Formation (Voss, 2002) in the Salar de Antofalla in north-west Argentina probably commenced accumulation in the latest Oligocene, but underwent their most important phase of accumulation during the Lower to Middle Miocene. These units are between 100 m and 150 m thick and record the interaction between aeolian and fluvial systems.

One of the best-described examples of an aeolian system developed in the Andean foreland is the Lower Miocene Vallecito Formation (Tripaldi & Limarino, 2005). This unit attains a maximum thickness of 1200 m and comprises a complex facies arrangement that records the interaction of dunes, aeolian sandsheets and wet interdunes that interact with fluvial and lacustrine systems. The succession records a large aeolian system developed as the first syn-orogenic fill of the Andean foreland in this part of north-western Argentina (Tripaldi & Limarino, 2005). Another Lower Miocene unit with similar characteristics is the Pachaco Formation in the Precordillera of San Juan in western Argentina. The middle member of this formation is 700 m thick and records the accumulation of a large dunefield dominated by barchan and seif dunes and draas (Milana *et al.*, 1993). The Angastaco Formation is related to an aeolian dunefield associated with fluvial systems

that developed in the Lower Miocene of north-western Argentina (Do Campo *et al.*, 2010). Both the Mariño Formation in the Precordillera of Mendoza (Irigoyen *et al.*, 2000) and the Santo Domingo Member of the El Durazno Formation in the Sierra de Famatina (Dávila & Astini, 2003) also record synorogenic aeolian systems associated with the development of the Andean foreland during the Middle Miocene. In southern Patagonia, aeolian deposits have also been described as part of the Lower to Middle Miocene Santa Cruz Formation (Pinturas Formation of Bown & Larriestra, 1990), and in distal portions of the Andean Foreland and in the passive South American margin, sandy aeolian facies have been described as part of the Río Negro Formation associated with a marine transgression from the Atlantic (Zavala & Freije, 2001).

Most of the Neogene aeolian systems of Africa are closely related to Quaternary systems and they record early accumulation conditions, with some differences due to changes in climate over the past 10 Myr (Fig. 8B). The oldest record is the Middle Miocene Tsondeb Sandstone Formation in Namibia which comprises a succession up to 220 m thick of cross-stratified and massive sandstones with the local development of pedogenic carbonates and palaeosols (Ward, 1988; Kocurek *et al.*, 1999; Ségalen *et al.*, 2004). This unit is interpreted as a proto-Namib sand sea that was influenced by winds that blew from the south/south-west, as today, but which developed under more humid conditions than those experienced today (Kocurek *et al.*, 1999). The succession comprises two sequences, each separated by a stabilization surface; deposits record the preservation of north-trending linear dunes that gradually undertook a lateral component of migration to the east. These linear forms supported superimposed dunes, similar to the large linear bedforms of the present-day Namib Desert, mainly as a consequence of a sustained wind regime that has been established since the Miocene (Ségalen *et al.*, 2004). Elsewhere in the Namib, up to four aeolian sequences are recognized (Senut *et al.*, 1995), comprising the deposits of star, linear and transverse dunes (Ségalen *et al.*, 2004).

Miocene aeolian deposits have been described in the Chad Basin and these are associated with the early hominid specimens in the Toros-Menalla site 266, in northern Chad, central Africa (Schuster *et al.*, 2002; Vignaud *et al.*,

2002). Accumulation of these aeolian successions was related to the early development of the Sahara (Schuster *et al.*, 2002; Vignaud *et al.*, 2002), although the relevance of this finding and its implication for pre-Quaternary desert development has been disputed (Swezey, 2006).

Upper Miocene to Lower Pliocene aeolian deposits are also present in the Western Cape of South Africa. In this area, the Prospect Hill Formation is composed of calcarenites with shell fragments that record the development of a coastal dune system overlying sandy beach deposits (Franceschini & Compton, 2004).

The Garet Uedda Formation (or Members U and V of the Sahabi Formation) in Libya (Fig. 8C) has been described as Upper Pliocene in age and an aeolian origin has been proposed for this 25 m thick sequence of quartzitic sands interbedded with sandy shales (Tawadros, 2012). Some relic forms of the Kalahari Desert might be as old as Upper Pliocene and they record the aeolian reworking of fluvial sands (Lancaster, 2000; Haddon & McCarthy, 2005). These units have different formal names (Gordonia Formation, Kalahari Sand, Batéké Sands and Zambezi Formation) and they can be correlated from Zaire and south-western Angola in the north to Botswana, Namibia and northern South Africa in the south. Deposits of these successions are mainly unconsolidated sand and their distribution suggests a larger sand sea than the Quaternary Kalahari Desert, giving rise to the largest sand body on Earth (the 'Mega Kalahari') that covered over 2.5 million km² (Thomas & Shaw, 1990).

Neogene aeolian activity in Asia was dominated by the accumulation of thick sequences of fine-grained deposits (loess) in the Loess and Tibetan Plateaus and aeolian deposits interbedded with alluvial deposits (e.g. Zheng *et al.*, 2003). Sandstone aeolian accumulations are not common and only one example has been identified in the Middle East: the Shuwaihat Formation, a Middle Miocene unit exposed on a 16 m high cliff near Abu Dhabi in the United Arab Emirates (UAE; Whybrow *et al.*, 1999), interpreted as the record of the interaction between transverse and barchanoid dunes and a continental sabkha (Bristow, 1999).

Quaternary aeolian sand systems

Sand seas and dunefields occurring today developed during the Quaternary Era, during which

significant changes in climate and sea-level related to glacial–interglacial cyclicality affected the supply, availability and mobility of sediment. Their accumulation and present configuration therefore reflect the legacy of these changes, in addition to contemporary processes. Quaternary aeolian sand systems occur on all continents and at all latitudes, with major systems located between 45 degrees north and south (Fig. 9), and with other smaller systems occurring at higher latitudes, including in the Arctic and Antarctica. Quaternary aeolian systems are here sub-divided into those located inland and those occurring along the coast.

Inland sand seas and dunefields (Table S5)

Inland dune systems occur widely, with a concentration in low to mid-latitude arid regions of the Northern Hemisphere (35–50°N), especially in the arid regions of central Asia, on the semi-arid Great Plains of North America and in low-latitude desert areas of Africa, Arabia and Australia (15–30°N and 15–30°S). Their geological setting varies, with many sand seas in Africa, Australia and Arabia occurring in cratonic basins; central Asian and South American sand seas, by contrast, are located mostly in foreland basins (Fig. 10). Dune types also vary, with linear dunes comprising *ca* 50% of all dunes and

dominating in many areas of the Sahara, south-east Arabia, Australia and southern Africa (e.g. Pye & Tsoar, 1990; Lancaster, 1999). Crescentic dunes comprise 40% of dunes and dominate sand seas in the northern Sahara, many parts of Arabia, and parts of central Asia and China. Star dunes comprise *ca* 8% of dunes in low-latitude inland sand seas, mainly in areas where topography creates complex wind regimes.

Wet systems form inland where the depositional surface intersects local perched or regional groundwater tables. Good examples of wet systems are the White Sands dunefield in New Mexico (Fig. 11; Kocurek *et al.*, 2007) and the Liwa area of the United Arab Emirates (Glennie, 2005; Stokes & Bray, 2005). Changes in sea-level, climate and/or vertical crustal movements that affect the groundwater table may result in a system changing over time from wet to dry or vice versa, as in the Wahiba Sands of Oman (Radies *et al.*, 2004). Likewise, spatial changes in groundwater levels result in some parts of the system being dry and others wet, as in the Rub'al Khali sand sea of Saudi Arabia (Glennie, 1970; Al-Masrahy & Mountney, 2013).

It appears that the majority of modern and Quaternary aeolian sand systems operate as dry systems in which the water table is significantly below the depositional surface, such that it has




Fig. 9. Location of major low-latitude and mid-latitude inland sand seas and dunefields, as well as coastal carbonate aeolianite deposits. After Sun & Muhs (2007).




Fig. 10. Tectonic setting of major inland sand seas and dunefields. Tectonic data from Fugro Tellus Sedimentary Basins of the World Map <http://www.datapages.com/AssociatedWebsites/GISOpenFiles/FugroTellusSedimentaryBasinsoftheWorldMap.aspx>. See also Table S5.

no effect on the dynamics of the dune system. The major controls are therefore sand supply, its availability for transport and mobility (magnitude and frequency of winds capable of transporting sand). The Namib Sand Sea (Fig. 12) is a good and comprehensively studied example of a dry system, sourced principally by sand from the interior of southern Africa via the Orange River (Garzanti *et al.*, 2012). Its accumulation is interpreted as the product of regional changes in wind regime, which result in a reduction of transport rates in the direction of transport, leading to deposition of sand by bedform climbing and dune growth. Estimates of angles of bedform climb made by Lancaster (1989) range from 0.003° for the linear dunes to 0.03 to 0.16° for crescentic dunes in the southern part of the sand sea. There is a clear spatial pattern of sand accumulation in central areas of the sand sea, represented by the equivalent or spread out sand thickness of complex linear dunes that reach 150 to 200 m in height (Lancaster, 1989; Bullard *et al.*, 2011) (Fig. 12). Isotopic and sediment budget estimates for the age of the sand sea converge at around one million years (Vermeesch *et al.*, 2010), but many of the dunes are relatively young, with optically stimulated luminescence (OSL) ages of 17 to 24 ka for compound linear dunes in the southern sand sea (Bubbenzer

et al., 2007) and <6 ka for linear dunes on the northern margin of the sand sea (Bristow *et al.*, 2007). Although there is evidence for episodic dune accumulation in the Namib, it appears that the persistence of arid to hyper-arid conditions throughout the Quaternary has promoted development of a very well-organized dune system.

Stabilized dune systems are those in which vegetation or some other factor (for example, permafrost and pedogenesis) periodically or continually stabilizes the substrate while the system remains active overall. Large-scale stabilized systems include the Thar Desert sand sea in India, much of the South-western Kalahari, the Simpson-Strzelecki in Australia (Fig. 13) and parts of the Negev-Sinai sand sea (Egypt and Israel). Characteristic dune forms of these systems are vegetated linear dunes and parabolic dunes. Quaternary climatic changes also resulted in some dune systems becoming stabilized at times of increased precipitation, as in the early Holocene, when well-developed soils developed on the dunes of the southern Sahara (Kocurek *et al.*, 1991; Lancaster *et al.*, 2002). Such episodes of soil formation are important for preservation of older episodes of dune accumulation.

Many studies have indicated that dunes in arid Australia have accumulated episodically and some are of great age (up to one million




Fig. 11. Example of a wet aeolian system: White Sands, New Mexico, USA: (A) cross-section of dunefield showing relations between aeolian and lacustrine sediments (after Kocurek *et al.*, 2007); (B) 3 to 5 m high crescentic ridges migrating across the surface of older crescentic dune deposits shown by exposed cross-bedding; (C) trench in an interdune area showing cross-bedded dune structures. Machete is *ca* 0.6 m long.

years) (Fujioka *et al.*, 2009). In many locations, the cores of linear dunes may exceed 380 ka in age, with multiple late Pleistocene and Holocene accumulation episodes (Fitzsimmons *et al.*, 2007; Lomax *et al.*, 2011). These episodes of dune growth have taken place without complete reworking of the dunes, in large part because of pedogenic alteration and stabilization of the deposits of older dune accumulation episodes by aeolian addition of clay and carbonate derived from nearby alluvial and lacustrine environments (Cohen *et al.*, 2010; Hesse, 2011). Successive episodes of stabilization, reworking

and dune growth result in an accretionary structure for the dunes, often associated with lateral migration in addition to dune extension (Rubin, 1990). Similar structures have been observed in stabilized (vegetated) linear dunes in other regions (e.g. Roskin *et al.*, 2011; Telfer, 2011).

Coastal systems (Table S5)

Coastal dunefields may be divided into those composed of siliciclastic material and those composed of carbonate (or mixed carbonate and siliciclastic) material. The former type typically develops along humid-type, mid-high latitude





Fig. 12. Example of a dry aeolian system: Namib Sand Sea, Namibia. Cross-section of central sand sea at 24.2°S; elevation data from ASTER GDEM; extent of Tsondab Sandstone Formation after Ward (1988). Landsat image of the area for comparison.

coasts, whereas the carbonate-rich dunes/dunefields form along arid to semi-arid, mid-low latitude, coasts bordering productive carbonate platforms.

Dune types vary and their size and morphology is dependent on a number of factors including vegetation cover, sand supply, wind regime and coastal setting. Sand blown off of the beach typically forms partly vegetated and fixed foredunes. Aeolian erosion of the foredunes can lead to the formation of blowouts and parabolic

dunes and/or transgressive dunefields (Hesp, 1999). Along cliffed coasts, special dune types including echo dunes and climbing dunes can develop (e.g. Clemmensen *et al.*, 1997).

Most coastal dunefields in North-west Europe can be classified as wet because the groundwater table typically is close to the surface. Depth to the groundwater table is, in many cases, linked to sea-level, especially in subsiding coastal basins (Kocurek *et al.*, 2001; Mountney & Russell, 2009). In other examples, especially in sys-




Fig. 13. Example of a stabilizing aeolian system: Strzelecki Desert, Australia. Along-dune profile showing multiple episodes of dune accumulation spanning the past 120 ka. From Cohen *et al.* (2010). 'II' to 'VII' – Marine isotope stages.

tems developed on uplifting coastal areas like the northern part of Denmark, dune dynamics are influenced by high precipitation rates and the formation of perched groundwater tables (Pedersen & Clemmensen, 2005; Clemmensen *et al.*, 2009). Dunefields develop both on retreating and prograding coasts. Both types may share many sedimentary characteristics, but prograding systems tend to develop successive lines of stabilized foredune ridges (e.g. Bristow & Pucillo, 2006; Madsen *et al.*, 2007; Reimann *et al.*, 2011), whereas retreating systems more commonly experience phases of transgressive dune formation in the form of inland migrating parabolic dunes (Clemmensen *et al.*, 2001a; Pedersen & Clemmensen 2005).

The Lodbjerg and Hvidbjerg coastal dunefields provide examples of wet-stabilized siliciclastic systems developed on a retreating coast (Figs 14 and 15). These two systems form part of an almost unbroken belt of coastal dunefields that flank the North Sea coast of Jutland, Denmark (Pedersen & Clemmensen, 2005; Clemmensen *et al.*, 2009). Luminescence dating of the sand units and radiocarbon dating of the peaty palaeosols have made it possible to establish a detailed chronology of dunefield evolution (Fig. 16). Episodes of transgressive dune formation that occurred around 2200 BC, 800 BC, 100 AD, 1050 to 1200 AD, and between 1550 and 1650 AD were linked to periods of increased storminess (cool, wet summers), whereas stabilization took place during periods of decreased storminess (Clemmensen *et al.*, 2009). The series of ages obtained by Clemmensen *et al.* (2001a) indicate accumulation of around 10 m of aeolian sand (below the present groundwater table) since 2200 BC, at an average rate of 2.4 mm yr^{-1} .

Swina barrier coastal dune system is another example of a wet-stabilized siliciclastic system

developed on a prograding shoreline. The Swina barrier is situated in north-west Poland along the southern part of the Baltic Sea; the dunefield is developed on top of two sandy spits that have formed between Pleistocene headlands (Reimann *et al.*, 2011). Spit formation and shoreline progradation have taken place during the past 6.6 kyr. The coast now forms a smooth and curved shoreline segment and is still prograding. The Swina barrier system is sourced by sand eroded from nearby headlands. Luminescence dating of the dunes indicates six hiatuses in foredune building, at 2100 BC, 900 BC, 200 BC, 200 AD, 600 AD, 1000 AD and 1600 AD. It is concluded that most of these phases of foredune erosion and instability were caused by climatic shifts to a cooler and windier climate. The transgressive dune formation *ca* 1600 AD was linked to increased storminess during the 'Little Ice Age' and this episode of dune formation seems to be contemporaneous with phases of increased aeolian activity in other dune systems in North-west Europe (e.g. Clemmensen & Murray, 2006; Clarke & Rendell, 2009; Clemmensen *et al.*, 2009).

Carbonate-rich aeolian systems are commonly developed in mid-latitude and low-latitude, arid and semi-arid climate belts; these aeolian systems occur in a variety of settings including lowland and cliffed coasts (e.g. Brooke, 2001; Frébourg *et al.*, 2008). Due to the climatic setting of these systems, they are most logically classified as dry. The carbonate sand is lithified soon after deposition (Guern & Davaud, 2005), thereby forming one mechanism of preservation that is poorly known from siliciclastic systems. These lithified carbonate-rich aeolian deposits are termed aeolianites (Brooke, 2001).

Particularly well-developed carbonate aeolian systems occur in the Western Mediterranean region (Clemmensen *et al.*, 1997; Fornós *et al.*,



Fig. 14. Ground-penetrating radar mapping of sedimentary architecture; coastal dunefield at Lodbjerg, Denmark (Clemmensen *et al.*, 2001a). The surface of the modern dunefield is situated *ca* 15 m above sea-level, and the dunefield is truncated by a coastal cliff towards the North Sea. Partly active cliff-top dunes are developed along the cliff. Vehicle for scale is *ca* 5 m long.




Fig. 15. Coastal cliff section at Lodbjerg, Denmark. A Wechselian till at the base of the section is overlain by 15 to 20 m of aeolian sand with peaty soils (dark horizons). Terminology of aeolian events after Clemmensen *et al.* (2009).

2009; Andreucci *et al.*, 2010a). Quaternary successions with carbonate-rich aeolian sand units crop out quasi-continuously along the north-west coast of Sardinia near the town of Alghero (Andreucci *et al.*, 2010a,b, 2014). The aeolian units, which are lithified, occur along a cliffed coast and can be subdivided into cliff-front dune accumulations and valley-head sand ramps (Andreucci *et al.*, 2010a; Fig. 17). Note also other major Quaternary aeolianites in South Africa (e.g. Roberts, 2008).

AEOLIAN RESEARCH: THE WAY FORWARD AND FUTURE RESEARCH PROSPECTS

Aeolian facies and sequence stratigraphic models: a useful approach to capturing complexity in aeolian successions?

Relating preserved aeolian stratigraphy to original bedform morphology and behaviour

Although it is now possible to effectively describe in detail both: (i) the morphological characteristics of modern bedforms and larger dunefields; and (ii) the geometry of architectural elements of preserved aeolian successions, notably by using the forward stratigraphic modelling techniques developed by Rubin (1987) and Rubin & Carter (2006), several problems remain regarding how to relate ancient preserved sets of aeolian strata to the morphology and migratory behaviour of the original bedforms. In particular,




Fig. 16. Simplified aeolian stratigraphy, Lodbjerg coastal dunefield. Note that most previous events of aeolian sand movement and transgressive dune formation led to the accumulation of aeolian sand-sheet deposits. Cross-bedded strata of inland migrating parabolic dunes are only preserved locally. The modern cliff-top dunes have eroded part of the underlying aeolian succession. Radiocarbon dating of the peaty soils and optically stimulated luminescence dating of the aeolian sand indicate that major periods of aeolian sand movement and transgressive dune formation were initiated at *ca* 2200 BC, 700 to 800 BC and AD 1100 to 1200. The cliff-top dunes have formed since about AD 1900. Terminology of aeolian events after Clemmensen *et al.* (2009).

problems remain in cases where large compound and complex morphological dune types have accumulated in desert basins in which the rate of accommodation generation has been highly variable over time or space, for example in response to spatially and temporally variable synsedimentary tectonic activity (e.g. Rodríguez-López *et al.*, 2013).

The accumulated sedimentary record of most modern inland dunefields remains largely unknown, with only fragmentary glimpses of aeolian sedimentary architectures having so far

been revealed from modern aeolian systems via techniques such as trenching (e.g. McKee, 1966) and ground-penetrating radar (GPR) studies (e.g. Bristow *et al.*, 2000a,b). Conversely, relatively few ancient aeolian successions are known that preserve, intact, the original morphologies of the bedforms that gave rise to the architecturally complex set and cosets of aeolian cross-bedding that dominate the ancient sedimentary record (e.g. Mountney *et al.*, 1999a,b). Thus, the development of aeolian facies models that are used to relate modern dunes and dunefields to accumulated deposits remains problematic in terms of how best to interpret the preserved sedimentary architectures of ancient aeolian deposits.

Despite great progress in understanding of the geomorphology of Quaternary inland sand seas, there are few data relating to their stratigraphic and sedimentological record. In many areas, Quaternary sand seas and dunefields have not left a significant accumulation and the bedforms present in many modern dunefields are known to be partially or completely legacy landforms inherited from Last Glacial Maximum (LGM) times (e.g. Lancaster *et al.*, 2002); thus, such forms do not necessarily reflect the currently prevailing climatic and sediment supply conditions. Elsewhere, information from the subsurface (for example, GPR data, cores and well logs) either does not exist or is proprietary. The costs and logistics of acquiring such data sets are often prohibitive but, where they have been developed, the understanding of the sedimentary record of inland sand seas has been transformed and relations between contemporary processes and the sedimentary record have been established (Bristow *et al.*, 2000a,b, 2005, 2007; Radies *et al.*, 2004; Stokes & Bray, 2005; Kocurek *et al.*, 2007; Derickson *et al.*, 2008).

There is a clear need to advance knowledge in this area by a coordinated programme of research that seeks to link dune processes to the sedimentary record. One valuable step in this direction is the development of a methodology for reconstructing wind direction, wind speed and duration of wind events from aeolian cross-strata (e.g. Eastwood *et al.*, 2012). The stratigraphic record of Quaternary (Holocene) coastal dunefields is better documented because such systems have been investigated by GPR mapping, outcrop and core studies, and placed in a chronological framework by luminescence and radiocarbon dating (e.g. Clemmensen *et al.*, 2001a; Pedersen & Clemmensen 2005), but in these systems, there is still a need to link aeolian processes more closely



Fig. 17. Valley-head sand ramp accumulation, north-west Sardinia (Andreucci *et al.*, 2010a). The sand ramp is composed of marine bioclastic-rich sand that was blown inland by dominant north-westerly winds *ca* 75 ka. Several episodes of ascending dune accumulation are revealed in the cliff section. Person for scale is *ca* 1.8 m tall.

to the sedimentary record. Sedimentary units with flat-bedded strata form a large portion of the stratigraphic record of these Holocene coastal systems (Clemmensen *et al.*, 2001a), but it is not yet clear whether these facies represent the trailing edges of inland migrating parabolic dunes or featureless sand plains formed in between the parabolic dunes.

The sedimentary structures of modern dunes need to be described more systematically in the literature. This includes the internal structures of fixed linear dune ridges, as well as those of many transgressive (actively migrating) dunes including the parabolic dunes of coastal areas. Studies of the internal sedimentary structures of dunes can be conducted from some natural exposures, including blow-outs, but the most systematic observation is provided from GPR

mapping of the dune bodies (e.g. Bristow *et al.*, 1996, 2000a,b; Clemmensen *et al.*, 2007; Tamura *et al.*, 2011). Such studies of internal structures should be accompanied by geomorphological studies, map studies of dune evolution and data on wind patterns. Bristow & Hill (1998) attempted to reconstruct changes in dune morphodynamics and ancient wind conditions from preserved sets of cross-stratification in the Miocene Shuwaihat Formation, using a reconstruction of localized wind patterns in relation to position on dune bedform slopes. This detailed reconstruction differs in scale to the more commonly attempted reconstructions of regional palaeowinds.

To date, relatively few pre-Quaternary coastal dune deposits (particularly in siliciclastic environments) have been described in the literature. Many ancient coastal dune deposits may have been overlooked and more studies of the internal structures of such modern dunes are needed to create a database of varied examples of sufficient size that will enable facies models for coastal aeolian dune systems to be ‘distilled’, and from which criteria for the recognition of characteristic stratigraphic architectures indicative of different coastal dune types can be established and applied to the stratigraphic record.

Recognition of the deposits of different aeolian bedform types in the preserved record

One fundamental problem that remains in relation to the interpretation of original dune type from the stratigraphic architecture preserved in the rock record is that the majority of reconstructions envisage transverse (and related barchanoid) dune types, whereas successions representing linear and star (and other) bedform types remain apparently under-recognized, despite such bedforms forming the majority of bedforms in modern dunefields. Given that there is no reason to suspect that linear and star bedforms were less abundant in the geological past, this suggests that current models and methods for the recognition and reconstruction of primary bedform type from preserved sedimentary architectures are not sufficiently refined to reliably distinguish bedform type. The reasons for these limitations in current facies models probably involve one or more of the complex factors considered below.

Linear bedforms, which make up *ca* 50% of the dunes in the world’s modern dunefields, principally transport sediment in an orientation close to parallel to the trend of their

crestlines; however, Bristow *et al.* (2000a, 2007) used GPR techniques to convincingly demonstrate that a minor component of transverse motion in such bedforms favours the preferential preservation of cross-strata that dip in the direction of that transverse component of migration. Thus, such linear forms tend to produce cross-strata that dip towards an azimuth that is at a high orientation to the migration direction of the primary bedform. The accumulation of sets of such cross-strata is therefore very difficult to distinguish from the deposits of transverse bedforms. This has significant implications for the reconstruction of palaeowind directions based on analysis of dip-azimuth data determined from preserved foresets, which, for linear bedforms, could be in orientations oblique to the dominant wind direction (see Scherer, 2000). Although recognition of linear draas is not an easy task, sedimentological and architectural features have been proposed as useful tools for their recognition in the sedimentary record (see Scherer, 2000; Rodríguez-López, *et al.*, 2008).

Many linear and star bedforms in present-day dunefields are isolated from neighbouring bedforms by extensive interdune flats and this suggests that such dunefields are not necessarily currently undergoing active accumulation, even in cases where the bedforms might be undergoing active construction (see Kocurek, 1999). Indeed, many large-scale examples of such bedforms are currently partially or fully stabilized by vegetation, by surface crusts or because the present-day wind regime is not capable of inducing the migration of such large forms (Bristow & Mountney, 2013). As such, linear and star forms (and their deposits) might have a reduced accumulation and, by implication, preservation potential such that successions representing the preserved deposits of such bedforms are indeed rare in the ancient stratigraphic record.

Draa architecture and apparent scarcity in the sedimentary record

Numerous facies models have been proposed with which to predict the likely arrangement of complex cosets of cross-strata arising from the accumulation of compound or complex draas representative of landforms comprising two or more scales of bedform that occur superimposed upon one another and which might undergo migration in different directions and styles, and at different rates (e.g. Brookfield, 1977; Rubin & Hunter, 1985; Rubin, 1987). Large draas are

ubiquitous in the central parts of most of the world's major desert dunefields where interdunes have been reduced to isolated depressions and neighbouring bedforms can be shown to be migrating over one another to leave an accumulation. From a conceptual standpoint, the preservation potential of such draa deposits ought to be relatively good where such bedforms accumulate in gradually subsiding basins or under the influence of a sustained episode of water-table rise. However, documented examples of draa deposits preserved in the ancient geological record are relatively scarce (e.g. Clemmensen, 1989; Chrintz & Clemmensen, 1993; Herries, 1993; Mountney, 2006a; Veiga & Spalletti, 2007; Langford *et al.*, 2008; Rodríguez-López *et al.*, 2008, 2012a).

Obvious research questions that remain to be addressed in relation to this issue are as follows: (i) are the preserved deposits of large draas actually relatively common in the ancient aeolian record but under-recognized because detailed studies focused specifically on their reconstruction have not yet been undertaken, either because outcropping examples have not been recognized or because the style of outcrop is not appropriate for their convincing three-dimensional reconstruction; and (ii) is the preservation potential of large draas low such that well-preserved examples are actually rather rare in the ancient aeolian record? In relation to this second point, how might large draa bedforms become preserved? In situations where such bedforms migrate over one another, angles of climb would probably be very low and, as such, only the very lowermost flanks of these large bedforms would accumulate. Such deposits would tend to be dominated by sets of low-angle inclined cross-strata dominated by wind-ripple deposits that represent only the lower plinths of large bedforms. Reconstruction of the original bedform morphology from this fragmentary record would be problematic. The potential for the preservation of such large bedforms essentially *intact* is low and would probably require exceptional circumstances, such as inundation by flood basalts (e.g. Jerram *et al.*, 1999a,b; Mountney *et al.*, 1999a,b) or important transgressive events (Strömbäck *et al.*, 2005; Rodríguez-López *et al.*, 2012a).

Mechanisms of accumulation and preservation: alternatives to bedform climbing
Many past and current facies models that attempt to account for the preserved sedimen-

tary architecture of aeolian successions invoke bedform climbing as a mechanism to explain the accumulation of thick preserved successions of aeolian dune and interdune strata. In its most basic form, the bedform-climb model envisages a train of contemporaneously active bedforms separated by interdunes that undertake gradual and progressive accumulation as they migrate over time, thereby accumulating a series of stacked sets of cross-bedded dune and interdune strata. Crucially, accumulation via such a mechanism requires a sustained episode of accumulation that is concurrent with ongoing bedform migration. Although bedform climbing can be convincingly shown to have been responsible for some preserved successions (e.g. Mountney & Thompson, 2002), others are more likely to have accumulated in response to punctuated accumulation whereby the preserved record represents numerous sequences, each bounded by supersurfaces that represent potentially protracted episodes of non-deposition (bypass) or erosion (deflation) (e.g. Loope, 1985). Although accumulation via bedform climbing and accumulation to form individual sequences separated by supersurfaces have traditionally been considered competing end-member models (e.g. Kocurek, 1991), several studies now convincingly demonstrate that both mechanisms for accumulation can occur and that transitions between accumulation via bedform climb and via sequence and supersurface generation can occur both spatially and temporally for some preserved aeolian successions (e.g. Mountney, 2012). Indeed, other mechanisms for the accumulation of aeolian successions are also possible (and arguably likely): the model of Paola & Borgman (1991) for the generation of sets of cross-strata in response to the migration of bedforms of pseudo-random height and scour depth and in the absence of net deposition (i.e. a zero angle of net climb) describes one such mechanism. However the long-term preservation potential of sets generated by such a mechanism remains uncertain because the passage of later bedforms would render the deposits prone to reworking in the absence of a component of accommodation generation.

Some research questions that remain to be answered include: (i) what proportion of the known aeolian preserved record arose via sustained episodes of bedform climb versus episodic sequence and supersurface generation, respectively; (ii) what specific set of controlling conditions allow for sustained bedform climb

versus what set of conditions allow for episodic sequence accumulation and preservation; and (iii) can sophisticated 4D models be developed with which to account for complex patterns of spatial and temporal transition between different mechanisms of accumulation and preservation and, if so, will such models be dependent on the development of high-resolution dating techniques, meaning that they will probably be best suited to the analysis of Quaternary aeolian successions?

Encapsulation of time in the aeolian stratigraphic record

It has long been recognized that the preserved sedimentary record is highly fragmentary (e.g. Barrel, 1917) such that the majority of geological time is encapsulated within unconformities. This notion is summarized succinctly by Ager (1993, p. 52) in the often-recounted statement that: “*the stratigraphic record is of one long gap with only very occasional sedimentation*”. Indeed, the aeolian record is likely to be especially fragmentary because both accumulation of aeolian deposits and also their long-term preservation demand a rather specific set of circumstances to conspire fortuitously (Kocurek, 1999), such that accumulation can take place at a time when accommodation space is being generated to enable preservation. Indeed, wind-blown sands deposited on thick continental crust are especially likely to be riddled with unconformities (e.g. Loope, 1985).

Several significant research questions remain to be addressed regarding the duration required to construct, accumulate and ultimately preserve aeolian system deposits. Specifically, how much time is represented by thick accumulations of aeolian strata as a fraction of the total time episode over which the succession accumulated? How much time is encapsulated within the unconformities (i.e. supersurfaces) present in successions of multiple stacked aeolian erg sequences and what are the fundamental allogenic controls that act at the basin scale to govern the timing of episodes erg construction and accumulation versus episodes of supersurface generation? (see Rodríguez-López *et al.*, 2013) Are episodes of aeolian construction and accumulation and episodes of supersurface formation gradual or abrupt, or does the time required for their development vary according to the underlying nature of the climatic and tectonic controls? The role of local synsedimentary tectonics (basin setting) and the differences between ergs

accumulated in different tectonic settings (for example, foreland basins versus rift basins) remain largely open research questions. How do rates of accommodation generation (via basin subsidence), uplift and denudation vary between desert basins and how might such rates be reconstructed from the preserved aeolian record? Given that the preserved stratigraphic record is generally accepted to be highly fragmentary (e.g. Loope, 1985), are specific episodes during which accumulation and preservation occurred necessarily representative of conditions that prevailed throughout the entire episode over which aeolian systems were active more generally but for which no record was preserved? Alternatively (and arguably, more likely), is the preserved aeolian stratigraphic record inherently biased such that certain types of aeolian dunefields have a greater chance of preservation because the climatic and tectonic settings in which they are constructed and accumulated favour preservation? What specific sets of circumstances are required to enable accumulation and preservation? Over what duration can an aeolian sequence accumulate given favourable conditions?

Attempts to address these research issues face a fundamental problem: the availability of appropriate dating techniques with which to establish the time periods over which accumulation and preservation occurred. Attempts to date aeolian sequences are best suited to studies of Quaternary successions for which techniques such as high-resolution OSL and radiocarbon dating can be applied; yet establishing the presence, lateral extent and significance of major supersurfaces of regional extent is best suited to ancient outcropping successions for which key stratal surfaces can be traced over distances of many hundreds to thousands of square kilometres (e.g. Havholm *et al.*, 1993; Havholm & Kocurek, 1994; Blakey *et al.* 1996).

An additional problem is that supersurfaces may not necessarily form isochronously across an entire basin; rather, their development may proceed in a progressive manner across the basin as the conditions responsible for their development gradually change over time and space (cf. Porter, 1986). Such a complex evolutionary history could mean that a single supersurface may potentially be time transgressive and may exhibit different characteristics across its area in response to spatial variation in the allocyclic

controls responsible for its development at a basin scale (Rodríguez-López *et al.*, 2013).

The role of the water table in enabling accumulation and preservation

Although many aeolian systems and preserved successions are considered to represent dry aeolian systems, a significant number of other examples reveal evidence to suggest that the water table (or its near-surface capillary fringe) played a significant role in assisting accumulation and enabling long-term preservation (Mountney, 2012; cf. Kocurek & Havholm, 1993). Although several well-known aeolian successions, including parts of the Permian Cedar Mesa Sandstone (Mountney & Jagger, 2004) and the Jurassic Entrada Sandstone (Crabaugh & Kocurek, 1993), can be shown to have accumulated via bedform migration and climb that was coincident with a gradual and progressive rise in the relative water table. It is clear that these successions are divided into a series of separate sequences bounded by supersurfaces. Several questions remain unresolved in relation to the role of the water table in enabling aeolian accumulation and preservation; for example, over what period can a relative rise in water-table level be sustained to enable wet system accumulation and how thick might the resulting preserved wet aeolian sequences be?

Can alternatives to the 'climbing wet system' model be developed to account for the origin of individual aeolian dune sets preserved between sets of interdune strata indicative of a wet or damp substrate? Might such single aeolian dune sets record the sporadic accumulation and preservation of thin aeolian sequences during episodes when controlling conditions conspired to fortuitously enable preservation? If so, might the intervening 'interdune' accumulations actually record protracted episodes during which accumulation of damp sand flats occurred at very slow rates (cf. Kocurek, 1981b; Crabaugh & Kocurek, 1993)?

Why are climbing wet aeolian systems only relatively rarely recognized in modern settings (e.g. Mountney & Russell, 2009), yet are increasingly interpreted from ancient successions? Might this reflect a bias that is inherent in the preserved stratigraphic record, whereby wet aeolian systems have an increased preservation potential because they progressively subside beneath a relative water table as they accumulate and this protects these deposits from later

deflation (cf. Stokes, 1968; Loope, 1985; Kocurek & Havholm, 1993)?

The role of groundwater level in preserving aeolian sand accumulations in coastal settings has been emphasized by many authors (e.g. Kocurek *et al.*, 2001; Clemmensen *et al.*, 2001b; Mountney & Russell, 2009) but few studies have discussed in detail what actually controls the groundwater level in coastal systems (e.g. Bristow *et al.*, 2000b). Sea-level is frequently cited as a dominant control on groundwater level (Havholm & Kocurek 1994; Kocurek *et al.*, 2001), but in many coastal dune systems in northern Denmark, for example, groundwater level is up to 15 m above sea-level (e.g. Clemmensen *et al.*, 2001a). Clearly, factors other than sea-level are controlling the groundwater level in these coastal systems.

The problem of dating aeolian sediments and rocks

Age control for episodes of dune construction, stability and reworking, as well as rates of dune accumulation for many Middle and Late Pleistocene and Holocene aeolian systems is being provided by luminescence dating techniques (e.g. Murray & Clemmensen, 2001; Lancaster, 2008; Singhvi & Porat, 2008; Fornós *et al.*, 2009; Andreucci *et al.*, 2010b; Reimann *et al.*, 2011; Thomas, 2013). In most studies, luminescence dating is based on quartz grains having typical grain-sizes between 0.18 mm and 0.25 mm. A luminescence age is derived by dividing the dose absorbed from ionizing radiation during burial by the dose rate; the latter originates from the natural radioactivity (U, Th, ^{40}K and decay products) in the sediments and, to some degree, with cosmic radiation.

Supplementary age control can be obtained using K-rich feldspar grains (Buylaert *et al.*, 2012), and although quartz optically stimulated luminescence usually is limited to the last *ca* 100 kyr, accurate ages up to 600 kyr can be obtained using feldspar infrared stimulated luminescence (Buylaert *et al.*, 2012). Additional age control is provided by radiocarbon dating of organic-rich soils, which are common components of many Holocene coastal aeolian successions (e.g. Clemmensen *et al.*, 2001a, 2009; Arbogast *et al.*, 2002), but are rare in low and mid-latitude desert dune systems. Holocene aeolian sand systems may contain artefacts, and archaeological studies of these items can add to the chronological understanding of the evolu-

tion of the system (e.g. Kocurek *et al.*, 1991; Liversage & Robinson, 1993; Roberts, 2008). Some Quaternary aeolian deposits interbedded with shell-bearing beach or shallow marine deposits have been dated by means of U-series analysis or by use of aminostratigraphy (e.g. Hearty *et al.*, 1986; Hillaire-Marcel *et al.*, 1996), and a few Quaternary aeolian systems have been dated by magnetostratigraphy and susceptibility stratigraphic analysis (e.g. Nielsen *et al.*, 2004).

Pre-Quaternary aeolian successions are hard to date precisely due to the lack of material that can be dated numerically. Aeolian successions that are associated with lava flows, however, can be placed in a chronological framework (for example, the Cretaceous Etjo Formation of Namibia; Mountney & Howell, 2000). Most siliclastic aeolian deposits contain no fossils that can be used to place them in a precise biostratigraphic framework. However, some rare exceptions include the Tertiary Tsondeb Sandstone of Namibia (Pickford & Senut, 1999). In contrast, carbonate aeolianites contain abundant microfossils that potentially could be useful in biostratigraphic analysis (see Babić *et al.*, 2013). However, the application of such dating techniques should be undertaken with care to eliminate the risk of possible error associated with aeolian reworking of older bioclasts which could be incorporated into more recent dune systems (see Teller *et al.*, 2000).

Aeolian systems interbedded with fossiliferous marine strata can be placed in a biostratigraphic framework and commonly also interpreted in a sequence or climate stratigraphic context (for example, the Late Carboniferous Hermosa Formation, Utah, USA, Atchley & Loope, 1993; the Pennsylvanian to Permian lower Cutler beds, Utah, USA, Jordan & Mountney, 2010, 2012). Additional methods that have been used in chronological studies of pre-Quaternary aeolian successions include combined palaeomagnetic and cyclostratigraphic analysis (for example, aeolian sandstones associated with lacustrine sediments in the Triassic of eastern USA and East Greenland; Kent & Tauxe, 2005).

Quantitative modelling of aeolian systems

Quantitative forward stratigraphic modelling provides a tool with which to demonstrate the link among: (i) aeolian bedform morphology; (ii) the processes operating on and between such

bedforms; (iii) the style of migratory bedform behaviour; and (iv) the sedimentary architecture of preserved aeolian successions, which is characterized by packages of lithofacies arranged into architectural elements with predictable geometries that are delineated by bounding surfaces of various types.

Building on the innovative geometrical modelling approach of Rubin (1987), there is a clear need to develop an effective approach to the modelling and characterization of the distribution of aeolian lithofacies within larger-scale architectural elements and establishment of a method to unequivocally demonstrate the primary processes responsible for the generation of such lithofacies distributions. Forward stratigraphic models for aeolian systems that account for both spatial and temporal variations in original bedform morphology, scale and style of migratory behaviour are needed. Such *dynamic* models (cf. Mountney, 2006b, 2012) will be useful to predict the 4D evolution of aeolian systems and the 3D distribution of resultant aeolian architectural elements, which will vary both spatially and vertically (as a function of temporal system evolution).

Another outstanding research issue is the establishment of a modelling technique with which to demonstrate the relation between the development of bounding surfaces due to autogenic processes, such as bedform migration and climb, and those that might be influenced by allogenic controls. For example, bedform climbing to generate an accumulation (which is generally considered to be an autogenic process) requires a sustained supply of sediment that is available for transport, and a wind that is capable of carrying that sediment to the site of accumulation (Kocurek & Lancaster, 1999); each of these controls will probably be influenced by allogenic factors, such as climate change, tectonism or relative sea-level change (in the case of near-coast aeolian systems). It is now evident that many processes once considered to be either allogenic or autogenic in origin are, in fact, more complex and reflect combined extrinsic and intrinsic controls (see Veiga *et al.*, 2005; Mountney, 2012; Rodríguez-López *et al.*, 2012a).

Forward stratigraphic modelling should aim to provide a quantitative approach to the 4D sequence stratigraphic modelling of aeolian successions, whereby the generation of aeolian sequences and their bounding supersurfaces (for example, deflation surfaces) can be accounted for in terms of formative processes. Coupled

with advances in the application of dating techniques to Quaternary aeolian successions, such modelling approaches could be used to demonstrate the longevity (or otherwise) of episodes of aeolian dunefield construction and accumulation versus episodes of destruction and super-surface development (cf. Loope, 1985; Kocurek & Havholm, 1993; Mountney, 2006a; Jordan & Mountney, 2012; Rodríguez-López *et al.*, 2012a, 2013).

Crucially, any adopted modelling approach must be undertaken in combination with detailed studies of outcropping successions, whereby meticulous recording of key stratal surfaces (such as supersurfaces of regional extent) using recently available digital technologies such as laser scanning (LiDAR) and photogrammetry will provide the hard data with which to constrain and justify modelled synthetic stratigraphies. One particularly important application of this combined and integrated outcrop analysis and conceptual modelling technique will be to test scenarios for which accumulation and preservation might take place in response to bedform climb, versus scenarios for which episodic and punctuated accumulation of aeolian sequences bounded by supersurfaces might occur.

Potential applications of new approaches to forward stratigraphic modelling of aeolian systems and their preserved successions have been applied to better understand aeolian hydrocarbon reservoirs such as the Permian Rotliegend Group of North Sea (Howell & Mountney, 1997), the Permian Unayzah Formation of Saudi Arabia (Melvin *et al.*, 2010) and the Jurassic Norphlet Sandstone of the Gulf of Mexico (Story, 1998; Ajdukiewicz *et al.*, 2010). Future aeolian modelling applications include: (i) the development of more realistic object-based reservoir models for the improved characterization of net and non-net aeolian reservoir intervals; (ii) the prediction of the likely distribution and lateral extent of low permeability barrier and baffles to flow, for example due to the presence of mudstone units in interdune elements, or calcite cements (e.g. Arribas *et al.*, 2012) (iii) the prediction of fluid-flow pathways (Lindquist, 1988; Garden *et al.*, 2005); and (iv) a technique for reconstructing 3D aeolian architecture from attributes present in 1D core samples (e.g. Romain & Mountney, 2014). Similar approaches could also be used for the characterization of water aquifers (e.g. Bloomfield *et al.*, 2006) and for the assessment of aeolian successions as potential sites for the

underground sequestration of CO₂ (Chiaramonte, 2008).

How can studies of Quaternary dunefields and sand seas inform understanding of ancient preserved aeolian successions in the rock record?

Preceding sections have highlighted the issues and challenges involved in understanding the preserved record of aeolian strata. In many cases, the record of Quaternary aeolian accumulation can provide data to inform these questions.

Relating preserved strata to original bedform morphology requires better documentation of the sedimentary structures and architecture of modern dunes of all types. Such documentation will also aid the recognition of different dune types in the sedimentary record and will address the question posed by Rubin & Hunter (1985): why are deposits of longitudinal (linear) dunes rarely recognized in the sedimentary record? Ground-penetrating radar can provide such data in an efficient manner, but attempts at 3D reconstructions of the sedimentary architecture of modern dunes are rare (see Bristow *et al.*, 2007, for an example). There exists a clear need for a comprehensive program of 3D GPR imaging of a representative selection of modern dunes to provide a 'library' of architectural styles for comparison with the rock record. Such studies will need to be guided by selection of appropriate spatial scales for acquisition of GPR data and make use of innovative software applications for visualization of these complex data sets.

The nature of the intervals of time represented by episodes of accumulation separated by bounding surfaces of different orders can now be addressed, at least for Quaternary successions, by high-resolution OSL dating of dunes and their accumulations and radiocarbon dating of associated organic-rich soils, coupled with stratigraphic information from exposures or GPR (e.g. Clemmensen *et al.*, 2001a,b; Pedersen & Clemmensen, 2005). A promising development is the modelling of dune accumulation using a probabilistic approach (Telfer *et al.*, 2010; Bailey & Thomas, 2013). The discontinuous nature of many dune accumulations is becoming very clear. Optically stimulated luminescence dating studies in several areas suggest that intervals of rapid accumulation of sand separated by long hiatuses is a common feature of dune accumulation (e.g. Bray and Stokes, 2003; Bristow *et al.*, 2007; Atkinson *et al.*, 2011; Telfer, 2011). However,

Leighton *et al.* (2013) have conducted a detailed sampling of aeolian sedimentary units in large linear dunes in the UAE, and conclude that stratigraphy alone is not sufficient at the studied sites to guide OSL sampling. For example, significant unconformities that represent a hiatus in dune accumulation are not distinguishable in the field from bounding surfaces representing more rapid changes in dune dynamics, whereas unconformities separating stratigraphic units of similar sedimentary properties do not always conform to the duration of associated hiatuses determined from OSL ages of the units.

On a broader spatial and temporal scale, a synthesis of multiple data sets to determine the boundary conditions of tectonics, topography and climate in which Quaternary aeolian sand systems have developed would provide a valuable baseline against which to evaluate ancient sand seas and dunefields. Such an approach requires a data intensive investigation that transcends the efforts made by Breed *et al.* (1979), and will require collection of data that describe aspects of spatially changing dune and interdune morphology and migratory behaviour across dunefields and their marginal areas (e.g. Al-Masrahy & Mountney, 2013). Coastal dunefields should be treated separately and dunefield evolution in particular related to coastal setting, sea-level and climate change. Similar approaches, most notably in relation to Distributive Fluvial Systems (DFS), have transformed our ability to understand these systems and have significantly altered perspectives on how and where these systems form and are preserved in the rock record (Hartley *et al.*, 2010; Weissmann *et al.*, 2010), leading to a new interest in the sedimentology of these features and their relations to groundwater and hydrocarbon reservoirs. The work of Livingstone *et al.* (2010) in Namibia is a promising start to this type of study.

What will be the ultimate preserved record of the great Last Glacial Maximum sand seas?

The period during the Last Glacial Maximum (19 to 26 ka) and immediately following it (11.5 to 19 ka) is characterized by widespread evidence of inland dune and draa construction. Preservation of the deposits of these landforms and their eventual incorporation into the rock record is, however, much less certain and requires that preservation space (*sensu* Kocurek,

1998) exists as a result of subsidence and burial, sea-level rise and/or a relative or absolute rise in the water table. There are few documented examples of any of these situations. Subsidence alone is usually insufficient to preserve aeolian deposits unless accompanied by a rise in water table that limits bedform migration and erosion. Such conditions may have existed locally during early Holocene wet phases in the Sahara and Arabia, but subsequent aridification has resulted in falling water tables and reworking of accumulations, as in the Liwa area of the UAE where Marine Isotope Stage 5 (MIS-5) and Holocene aeolian deposits dominate (Stokes & Bray, 2005). Similarly, during the early Holocene, dunes in the southern and western Sahara and parts of Arabia were stabilized by soil formation (Talbot, 1985) and this stabilized surface has survived in many places to the present day.

A promising environment for preservation of Last Glacial Maximum dune deposits may occur where dunes from inland migrated or extended onto the continental shelf during glacial periods of low sea-level and were subsequently preserved by shallow marine and/or coastal sabkha deposits during the early Holocene sea-level rise. Such scenarios exist in Mauritania (Lancaster *et al.*, 2002; Hanebuth *et al.*, 2013), in the Arabian Gulf (Shinn, 1973; Al-Hinai *et al.*, 1987), where crescentic dunes have built a clastic wedge offshore of Qatar, overlain by coastal sabkha deposits, and in north-western Australia, where linear dunes are preserved by estuarine muds deposited during the Holocene transgression (Jennings, 1975). Hanebuth *et al.* (2013) present a detailed analysis of the architecture of aeolian and shallow marine deposits off of northern Mauritania, which indicates preservation of deposits of an inland dune complex of MIS-4 age, but not of the MIS-2 (LGM) dunes, despite the intensity of this aeolian constructional phase. These authors hypothesize that LGM dune deposits are not preserved because they were not cemented prior to the sea-level rise. Alternatively, the position of major dune complexes may have shifted to the south between MIS-4 and MIS-2. In either case, the present study demonstrates the likely sporadic nature of preservation of inland dune deposits in the Quaternary sedimentary record.

During the Last Glacial Period global sea-level was low and during the Last Glacial Maximum sea-level had dropped to a level *ca* 120 m below

that of the present day. Luminescence dating indicates that these conditions of low sea-level frequently led to inland wind transport of bioclastic sand from exposed shelf areas (e.g. Zhou *et al.*, 1994; Preusser *et al.*, 2002; Radies *et al.*, 2004; Fornós *et al.*, 2009; Andreucci *et al.*, 2010a). Sand – commonly in the form of transgressive dunes – migrated far inland building large (coastal) dunefields such as the Wahiba Sands, Sultanate of Oman (Radies *et al.*, 2004), or more isolated aeolian accumulations such as cliff-front dune or ramp deposits on Mallorca and Sardinia, Western Mediterranean (e.g. Clemmensen *et al.*, 1997; Fornós *et al.*, 2009; Andreucci *et al.*, 2010a,b). Preservation of dune sand in the Wahiba Sands took place during a subsequent episode of increased wetness and elevated groundwater table (Radies *et al.*, 2004), whereas preservation of the cliff-front aeolian accumulations in the Western Mediterranean was linked to early cementation (Fornós *et al.*, 2009).

CONCLUSIONS

Wind has been an agent responsible for eroding, transporting and depositing sediments on Earth since at least 3.2 Ga. The study of the sedimentary record of aeolian deposits has allowed resolution of some fundamental questions regarding sedimentology and stratigraphic architecture. However, important questions regarding the complexity of the relative role of different allocyclic controls, preservation mechanisms and facies variability remain to be resolved. Multidisciplinary approaches are required to address a series of research problems of fundamental importance: has the terrestrial atmosphere had the same properties throughout Earth history, and how have they controlled aeolian systems development? Facies models of aeolian systems are currently too simplistic and do not account for the complexity inherent in facies patterns. The sedimentary record is likely to be fundamentally biased because preserved aeolian deposits represent only a fraction of the dunefields that have persisted throughout Earth history. Furthermore, common mechanisms for aeolian accumulation tend to preserve only the lowermost parts of dune aprons, such that attempts to reconstruct aeolian palaeoenvironments are based on data that reflects only a small part of an original dunefield system. The development of detailed facies and sequence stratigraphic models will require high-

resolution dating techniques, meaning that they will probably be best devised via the analysis of Quaternary aeolian successions. At the basin scale, the significance of synsedimentary tectonics has probably been understudied with respect to other allocyclic controls (for example, climate) which are known to influence erg architecture and supersurface generation. This study serves to outline the current limitation in current understanding of aeolian systems and their preserved successions: it is hoped that this discussion will stimulate thought and renewed activity in addressing shortcomings in existing knowledge.

ACKNOWLEDGEMENTS

We are grateful to the Chief Editor Stephen Rice and to the Associate Editor Charlie Bristow for their editorial work and useful comments and suggestions. We also thank Gary Kocurek, David Loope and Charlie Bristow for their constructive and valuable reviews and advice on how to improve the earlier versions of this manuscript. This paper is a contribution to the scientific project CGL2011-23717 (Ministerio de Ciencia e Innovación, MICINN, Gobierno de España). Mountney is grateful to Areva, BHP Billiton, ConocoPhillips, Nexen, Saudi Aramco, Shell, Tullow Oil and Woodside for funding his contribution to this research through their sponsorship of the Fluvial & Eolian Research Group at the University of Leeds. Rodríguez-López thanks Prof. Poppe L. de Boer for discussions on palaeogeographical distribution of ergs through Earth history. Clemmensen thanks the Carlsberg Foundation that supported his research on Quaternary aeolian systems. Lancaster's research has been sponsored in part by the National Science Foundation and the American Chemical Society Petroleum Research Fund. Veiga thanks the Consejo Nacional de Investigaciones Científicas y Técnicas (CONICET), the Agencia Nacional de Promoción Científica y Tecnológica (ANPCyT) and YPF S.A. for funding research on aeolian systems at the Centro de Investigaciones Geológicas.

REFERENCES

- Ager, D.V. (1993) *The Nature of the Stratigraphical Record*, 3rd edn. John Wiley & Sons, Chichester, New York, Brisbane, Toronto, Singapore, 151 pp.
- Ajdukiewicz, J.M., Nicholson, P.H. and Esch, W.L. (2010) Prediction of deep reservoir quality using early diagenetic process models in the Jurassic Norphlet Formation, Gulf of Mexico. *AAPG Bull.*, **94**, 1189–1227.
- Al-Hinai, K.G., Moore, J.M. and Bush, P.R. (1987) LANDSAT image enhancement study of possible submerged sand-dunes in the Arabian Gulf. *Int. J. Remote Sens.*, **8**, 251–258.
- Alimen, H. (1936) Étude sur le stampian du Bassin de Paris. *Soc. Géol. Fr. Mem.*, **31**, 1–304.
- Allen, P.A. and Marshall, J.E.A. (1981) Depositional environments and palynology of the Devonian south-east Shetland Basin. *Scot. J. Geol.*, **17**, 257–273.
- Al-Masrahy, M.A. and Mountney, N.P. (2013) Remote sensing of spatial variability in aeolian dune and interdune morphology in the Rub' Al-Khali, Saudi Arabia. *Aeolian Res.*, **11**, 155–170.
- de Almeida, R.P., Janikian, L., Fragoso-Cesar, A.R.S. and Marconato, A. (2009) Evolution of a rift basin dominated by subaerial deposits: the Guaritas Rift, Early Cambrian, Southern Brazil. *Sed. Geol.*, **217**, 30–51.
- Andreucci, S., Clemmensen, L.B. and Pascucci, V. (2010a) Transgressive dune formation along a cliffed coast at 75 ka in Sardinia, Western Mediterranean: a record of sea-level fall and increased windiness. *Terra Nova*, **22**, 424–433.
- Andreucci, S., Clemmensen, L.B., Murray, A.S. and Pascucci, V. (2010b) Middle to late Pleistocene coastal deposits of Alghero, northeast Sardinia (Italy). Chronology and evolution. *Quatern. Int.*, **222**, 3–16.
- Andreucci, S., Panzeri, L., Peter Martini, I., Maspero, F., Martini, M. and Pascucci, V. (2014) Evolution and architecture of a West Mediterranean Upper Pleistocene to Holocene coastal apron-fan system. *Sedimentology*, **61**, 333–361.
- Annels, A.E. (1989) Ore genesis in the Zambian Copperbelt, with particular reference to the northern sector of the Chambishi Basin. In: *Sediment-Hosted Stratiform Copper Deposits* (Eds R.W. Boyle, A.C. Brown, C.W. Jefferson, E.C. Jowett and R.V. Kirkham), *Geol. Assoc. Can. Spec. Pap.*, **36**, 427–452.
- Arbogast, A.F., Hansen, E.C. and Van Oort, V.M. (2002) Reconstructing the geomorphic evolution of large coastal dunes along the southeastern shore of Lake Michigan. *Geomorphology*, **46**, 241–255.
- Arribas, M.E., Rodríguez-López, J.P., Meléndez, N., Soria, A.R. and de Boer, P.L. (2012) Giant calcite concretions in aeolian dune sandstones; sedimentological and architectural controls on diagenetic heterogeneity, mid-Cretaceous Iberian Desert System. *Spain. Sed. Geol.*, **243–244**, 130–147.
- Atchley, S.C. and Loope, D.B. (1993) Low-stand aeolian influence on stratigraphic completeness: upper member of the Hermosa Formation (latest Carboniferous), southeast Utah, USA. *Int. Assoc. Sedimentol. Spec. Publ.*, **16**, 127–149.
- Atkinson, O.A.C., Thomas, D.S.G., Goudie, A.S. and Bailey, R.M. (2011) Late Quaternary chronology of major dune ridge development in the northeast Rub' al-Khali, United Arab Emirates. *Quatern. Res.*, **76**, 93–105.
- Babić, L., Zupanić, J., Vidović, J., Razum, I., Lužar-Oberiter, B. and Crnjaković, M. (2013) Preservation of hanging aeolian deposits in insular karst depressions: sediment sources and implications for the Pleistocene palaeogeography of the SE Adriatic archipelago. *Aeolian Res.*, **11**, 171–189.
- Barrel, J. (1917) Rhythms and the measurement of geologic time. *Geol. Soc. Am. Bull.*, **28**, 745–904.

- Bart, H.A.** (1977) Sedimentology of cross-stratified sandstones in Arikaree group, Miocene, southeastern Wyoming. *Sed. Geol.*, **19**, 165–184.
- Bailey, R.M. and Thomas, D.S.G.** (2013) A quantitative approach to understanding dated dune stratigraphies. *Earth Surf. Proc. Land.*, **39**, 614–631.
- Benan, C.A.A. and Kocurek, G.** (2000) Catastrophic flooding of an aeolian dune field: Jurassic Entrada and Todilto Formations, Ghost Ranch, New Mexico, USA. *Sedimentology*, **47**, 1069–1080.
- Benton, M., Cook, E. and Turner, P.** (2002) *Permian and Triassic Red Beds and the Penarth Group of Great Britain*, Geological Conservation Review Series, Vol. 24. Joint Nature Conservation Committee, Peterborough, 337 pp.
- Bigarella, J.J.** (1979) Cave sandstone and similar sandstones of Southern Africa (Triassic). In: *A Study of Global Sand Seas* (Ed. E.D. McKee). United States Department of the Interior, *U.S. Geol. Surv. Prof. Pap.*, **1052**, 236–238.
- Biswas, A.S.** (2005) Coarse aeolianites: sand sheets and zibar-interzibar facies from the Mesoproterozoic Cuddapah Basin, India. *Sed. Geol.*, **174**, 149–160.
- Blakey, R.C.** (1990) Stratigraphy and geologic history of Pennsylvanian and Permian rocks, Mogollon Rim region, central Arizona and vicinity. *Geol. Soc. Am. Bull.*, **102**, 1189–1217.
- Blakey, R.C.** (1996) Permian Eolian deposits, sequences, and sequence boundaries, Colorado Plateau. In: *Paleozoic Systems of the Rocky Mountain Region* (Eds M.W. Longman and M.D. Sonnenfeld), pp. 405–426. SEPM, Rocky Mountain Section, Denver, CO.
- Blakey, R.C. and Middleton, L.T.** (1983) Permian shoreline eolian complex in central Arizona: dune changes in response to cyclic sea-level changes. In: *Eolian Sediments and Processes* (Eds M.E. Brookfield and T.S. Ahlbrandt), *Dev. Sedimentol.*, **38**, 551–581.
- Blakey, R.C., Peterson, F. and Kocurek, G.** (1988) Synthesis of late Paleozoic and Mesozoic eolian deposits of the western interior of the United States. *Sed. Geol.*, **56**, 3–125.
- Blakey, R.C., Havholm, K.G. and Jones, L.S.** (1996) Stratigraphic analysis of eolian interactions with marine and fluvial deposits, Middle Jurassic Page Sandstone and Carmel Formation, Colorado Plateau, USA. *J. Sed. Res.*, **66**, 324–342.
- Bloomfield, J.P., Moreau, M.F. and Newell, A.J.** (2006) Characterization of permeability distributions in six lithofacies from the Helsby and Wilmslow sandstone formations of the Cheshire Basin, UK. In: *Fluid Flow and Solute Movement in Sandstones: The Onshore UK Permo-Triassic Red Bed Sequence* (Eds R.D. Barker and J.H. Tellam), *Geol. Soc. London Spec. Publ.*, **263**, 83–101.
- Bordy, E.M. and Catuneanu, O.** (2002) Sedimentology and palaeontology of upper Karoo aeolian strata (Early Jurassic) in the Tuli Basin, South Africa. *J. Afr. Earth Sci.*, **35**, 301–314.
- Bown, T.M. and Kraus, M.J.** (1988) Geology and paleoenvironment of the Oligocene Jebel Qatrani Formation and adjacent rocks, Fayum Depression, Egypt. *U.S. Geol. Surv. Prof. Pap.*, **1452**, 60.
- Bown, T.M. and Larriestra, C.N.** (1990) Sedimentary paleoenvironments of fossil platyrrhine localities, Miocene Pinturas Formation, Santa Cruz Province, Argentina. *J. Human Evol.*, **19**, 87–119.
- Bray, H.E. and Stokes, S.** (2003) Chronologies for Late Quaternary barchan dune reactivation in the southeastern Arabian Peninsula. *Quatern. Sci. Rev.*, **22**, 1027–1033.
- Breed, C.S., Fryberger, S.G., Andrews, S., McCauley, C., Lennartz, F., Geber, D. and Horstman, K.** (1979) Regional studies of sand seas using LANDSAT (ERTS) imagery. In: *A Study of Global Sand Seas* (Ed. E.D. McKee), *U.S. Geol. Surv. Prof. Pap.*, **1052**, 305–398.
- Bristow, C.S.** (1999) Aeolian and sabkha sediments in the Miocene Shuwaihat Formation, Emirate of Abu Dhabi, United Arab Emirates. In: *Fossil Vertebrates of Arabia* (Eds P.J. Whybrow and A. Hill), pp. 50–60. Yale University Press, New Haven, CT.
- Bristow, C.S. and Hill, N.** (1998) Dune morphology and palaeowinds from aeolian sandstones in the Miocene Shuwaihat Formation, Abu Dhabi, United Arab Emirates. In: *Quaternary Deserts and Climatic Change* (Eds A.S. Alsharhan and K.W. Glennie), pp. 553–564. AA Balkema, Rotterdam, the Netherlands.
- Bristow, C.S. and Pucillo, K.** (2006) Quantifying rates of coastal progradation from sediment volume using GPR and OSL: the Holocene fill of Guichen Bay, south-east South Australia. *Sedimentology*, **53**, 769–788.
- Bristow, C.S. and Mountney, N.P.** (2013) Aeolian landscapes, Aeolian stratigraphy. In: *Treatise on Geomorphology* (Ed. J. Shroder). Elsevier, San Diego, CA, **11**, 246–268.
- Bristow, C.S., Pugh, J. and Goodall, T.** (1996) Internal structure of aeolian dunes in Abu Dhabi determined using ground-penetrating radar. *Sedimentology*, **43**, 995–1003.
- Bristow, C.S., Bailey, S.D. and Lancaster, N.** (2000a) The sedimentary structure of linear sand dunes. *Nature*, **406**, 56–59.
- Bristow, C.S., Chroston, P.N. and Bailey, S.D.** (2000b) The structure and development of foredunes on a locally prograding coast: insights from ground-penetrating radar surveys, Norfolk, UK. *Sedimentology*, **47**, 923–944.
- Bristow, C.S., Lancaster, N. and Duller, G.A.T.** (2005) Combining ground penetrating radar surveys and optical dating to determine dune migration in Namibia. *J. Geol. Soc.*, **162**, 315–322.
- Bristow, C.S., Duller, G.A.T. and Lancaster, N.** (2007) Age and dynamics of linear dunes in the Namib Desert. *Geology*, **35**, 555–558.
- Brooke, B.** (2001) The distribution of carbonate eolianite. *Earth-Sci. Rev.*, **55**, 135–164.
- Brookfield, M.E.** (1977) The origin of bounding surfaces in ancient aeolian sandstones. *Sedimentology*, **24**, 303–332.
- Brookfield, M.E.** (2004) The enigma of fine-grained alluvial basin fills: the Permo-Triassic (Cumbrian Coastal and Sherwood Sandstone Groups) of the Solway Basin, NW England and SW Scotland. *Int. J. Earth Sci.*, **93**, 282–296.
- Brookfield, M.E.** (2008) Palaeoenvironments and palaeotectonics of the arid to hyperarid intracontinental latest Permian-Late Triassic Solway Basin (U.K.). *Sed. Geol.*, **210**, 27–47.
- Browne, M.A.E., Smith, R.A. and Aitken, A.M.** (2002) *Stratigraphical Framework for the Devonian (Old Red Sandstone) Rocks of Scotland South of a Line from Fort William to Aberdeen*. British Geological Survey Research Report, RR/01/04. 67 pp.
- Bubbenzer, O., Bödeker, O. and Besler, H.** (2007) A transcontinental comparison between the southern Namib Erg (Namibia) and the southern Great Sand Sea (Egypt). *Zbl. Geol. Paläontol. Teil 1, Heft 1/4*, 7–23.
- Buylaert, J.-P., Jain, M., Murray, A.S., Thomsen, K.J., Thiel, C. and Sohbaty, R.** (2012) A robust feldspar luminescence

- dating method for Middle and Late Pleistocene sediments. *Boreas*, **41**, 435–451.
- Bullard, J.E., White, K. and Livingstone, I.** (2011) Morphometric analysis of aeolian bedforms in the Namib Sand Sea using ASTER data. *Earth Surf. Proc. Land.*, **36**, 1534–1549.
- Cain, S.A. and Mountney, N.P.** (2009) Spatial and temporal evolution of a terminal fluvial fan system: the Permian Organ Rock Formation, South East Utah, USA. *Sedimentology*, **56**, 1774–1800.
- Cain, S.A. and Mountney, N.P.** (2011) Downstream changes and associated fluvial-aeolian interactions in an ancient terminal fluvial fan system: the Permian Organ Rock Formation, SE Utah. In: *From River to Rock Record* (Eds S. Davidson, S. Leleu and C. North), *SEPM Spec. Publ.*, **97**, 165–187.
- Caselli, A. and Limarino, C.O.** (2002) Sedimentología y evolución paleoambiental de la Formación Patquía (Pérmico) en el extremo sur de la sierra de Maz y cerro Bola, provincia de La Rioja, Argentina. *Rev. Asoc. Geol. Argentina*, **57**, 415–436.
- Cather, S.M., Connell, S.D., Chamberlin, R.M., McIntosh, W.C., Jones, G.E., Potochnik, A.R., Lucas, S.G. and Johnson, P.S.** (2008) The Chuska erg: paleogeomorphic and paleoclimatic implications of an Oligocene sand sea on the Colorado Plateau. *Geol. Soc. Am. Bull.*, **120**, 13–33.
- Chakraborty, T.** (1991) Sedimentology of a Proterozoic erg: the Venkatpur Sandstone, Pranhita-Godavari Valley, South-India. *Sedimentology*, **38**, 301–322.
- Chan, M.A.** (1989) Erg margin of the Permian White Rim Sandstone, SE Utah. *Sedimentology*, **36**, 235–251.
- Chan, M.A. and Kocurek, G.** (1988) Complexities in eolian and marine interactions: processes and eustatic controls on erg development. *Sed. Geol.*, **56**, 283–300.
- Chan, M.A. and Archer, A.W.** (2000) Cyclic eolian stratification on the Jurassic Navajo Sandstone, Zion National Park; periodicities and implications for paleoclimate. In: *Geology of Utah's Parks and Monuments* (Eds D.A. Sprinkel, T.C. Chidsey Jr. and P.B. Anderson), *Utah Geol. Assoc. Publ.*, **28**, 606–617.
- Chiaromonte, L.** (2008) Geomechanical Characterization and Reservoir Simulation of a CO₂ Sequestration Project in a Mature Oil Field, Teapot Dome, WY. PhD Thesis, Stanford University, 192 pp.
- Chrintz, T. and Clemmensen, L.B.** (1993) Draa reconstruction, the Permian Yellow Sands, northeast England. In: *Aeolian Sediments: Ancient and Modern* (Eds K. Pye and N. Lancaster), *Int. Assoc. Sedimentol. Spec. Publ.*, **16**, 151–161.
- Clarke, M.L. and Rendell, H.M.** (2009) The impact of North Atlantic storminess on western European coasts: a review. *Quatern. Int.*, **195**, 31–41.
- Clemmensen, L.B.** (1985) Desert sand plain and sabkha deposits from the Bunter Sandstone formation (L.Triassic) at the northern margin of the German Basin. *Geol. Rundsch.*, **74**, 519–536.
- Clemmensen, L.B.** (1987) Complex star dunes and associated aeolian bedforms, Hopeman Sandstone (Permo-Triassic), Moray Firth Basin, Scotland. In: *Desert Sediments: Ancient and Modern* (Eds L.E. Frostick and I. Reid), *Geol. Soc. London Spec. Publ.*, **35**, 213–231.
- Clemmensen, L.B.** (1988) Aeolian morphology preserved by lava cover, the Precambrian Mussartut member, Eriksfjord Formation, South Greenland. *Bull. Geol. Soc. Denmark*, **37**, 105–116.
- Clemmensen, L.B.** (1989) Preservation of interdraa and plinth deposits by the lateral migration of large linear draas (Lower Permian Yellow Sands, Northeast England). *Sed. Geol.*, **65**, 139–151.
- Clemmensen, L.B. and Abrahamsen, K.** (1983) Aeolian stratification and facies association in desert sediments, Arran Basin (Permian), Scotland. *Sedimentology*, **30**, 311–339.
- Clemmensen, L.B. and Blakey, R.C.** (1989) Erg deposits in the Lower Jurassic Wingate Sandstone, Northeastern Arizona: oblique dune sedimentation. *Sedimentology*, **36**, 449–470.
- Clemmensen, L.B. and Murray, A.** (2006) The termination of the last major phase of aeolian sand movement, coastal dunefields, Denmark. *Earth Surf. Proc. Land.*, **31**, 795–808.
- Clemmensen, L.B., Olsen, H. and Blakey, R.C.** (1989) Erg-margin deposits in the Lower Jurassic Moenave Formation and Wingate Sandstone, Southern Utah. *Geol. Soc. Am. Bull.*, **101**, 759–773.
- Clemmensen, L.B., Øxnevad, I.E.I. and de Boer, P.L.** (1994) Climatic controls on ancient desert sedimentation: some late Palaeozoic examples from NW Europe and the western interior of the USA. In: *Orbital Forcing and Cyclic Sequences* (Eds P.L. de Boer and D.G. Smith), *Int. Assoc. Sedimentol. Spec. Publ.*, **19**, 439–457.
- Clemmensen, L.B., Fornós, J.J. and Rodríguez-Perea, A.** (1997) Morphology and architecture of a late Pleistocene cliff-front dune, Mallorca, Western Mediterranean. *Terra Nova*, **9**, 251–254.
- Clemmensen, L.B., Pye, K., Murray, A. and Heinemeier, J.** (2001a) Sedimentology, stratigraphy and landscape evolution of a Holocene coastal dune system, Lodbjerg, NW Jutland, Denmark. *Sedimentology*, **48**, 3–27.
- Clemmensen, L.B., Andreasen, F., Heinemeier, J. and Murray, A.** (2001b) A Holocene coastal aeolian system, Vejers, Denmark: landscape evolution and sequence stratigraphy. *Terra Nova*, **13**, 129–134.
- Clemmensen, L.B., Bjørnsen, M., Murray, A. and Pedersen, K.** (2007) The formation of aeolian dunes on Anholt, Denmark since AD 1560: a record of deforestation and increased storminess. *Sed. Geol.*, **199**, 171–187.
- Clemmensen, L.B., Murray, A., Heinemeier, J. and De Jong, R.** (2009) The evolution of Holocene coastal dunefields, Jutland, Denmark: a record of climate change over the past 5000 years. *Geomorphology*, **105**, 303–313.
- Cohen, T.J., Nanson, G.C., Larsen, J.R., Jones, B.G., Price, D.M., Coleman, M. and Pietsch, T.J.** (2010) Late Quaternary aeolian and fluvial interactions on the Cooper Creek Fan and the association between linear and source-bordering dunes, Strzelecki Desert, Australia. *Quatern. Sci. Rev.*, **29**, 455–471.
- Cojan, I. and Thiry, M.** (1992) Seismically induced deformation structures in Oligocene shallow-marine and aeolian coastal sands (Paris Basin). *Tectonophysics*, **206**, 79–89.
- Correa, G.A., Carrecedo, M.L. and Gutierrez, P.R.** (2012) Paleambiente y paleontología de la Formación Andapaico (Paleozoico superior, Precordillera Central, Argentina). *Andean Geol.*, **39**, 22–52.
- Cowan, G. and Boycott-Brown, T.** (2003) The North Morecambe field, block 110/2a, East Irish Sea. *Geol. Soc. Am. Mem.*, **20**, 97–105.
- Crahaugh, M. and Kocurek, G.** (1993) Entrada Sandstone: an example of a wet aeolian system. In: *The Dynamics and*

- Environmental Context of Aeolian Sedimentary Systems* (Ed. K. Pye), *Geol. Soc. London Spec. Publ.*, **72**, 103–126.
- Crahaugh, M.** and **Kocurek, G.** (1998) Continental sequence stratigraphy of a wet aeolian system. In: *Relative Role of Eustasy, Climate and Tectonism in Continental Rocks* (Eds K. Stanley and P. McCabe), *SEPM Spec. Publ.*, **59**, 213–228.
- Dávila, F.M.** and **Astini, R.A.** (2003) Las eolianitas de la Sierra de Famatina (Argentina): interacción paleoclimatológica en el antepaís fragmentado andino central durante el Mioceno Medio? *Rev. Geol. Chile*, **30**, 187–204.
- De Ploey, J., Lepersonne, J.** and **Stoops, G.** (1968) *Sédimentologie et origine des sables de la série des Sables Ocre et de la série des « Grès Polymorphes » (système du Kalahari) au Congo Occidental*. Musée Royal de l'Afrique Centrale, Tervuren, Belgique, 72 pp.
- Derickson, D., Kocurek, G., Ewing, R.C.** and **Bristow, C.S.** (2008) Origin of a complex and spatially diverse dune-field pattern, Algodones, southeastern California. *Geomorphology*, **99**, 186–204.
- Deynoux, M., Kocurek, G.** and **Proust, J.N.** (1989) Late Proterozoic periglacial aeolian deposits on the West African Platform, Taoudeni Basin, western Mali. *Sedimentology*, **36**, 531–550.
- Dias, K.D.N.** and **Scherer, C.M.S.** (2008) Cross-bedding set thickness and stratigraphic architecture of aeolian systems: an example from the Upper Permian Piramboia Formation (Parana' Basin), southern Brazil. *J. S. Am. Earth Sci.*, **25**, 405–415.
- Dickinson, W.R.** and **Gehrels, G.E.** (2003) U-Pb ages of detrital zircons from Permian and Jurassic aeolian sandstones of the Colorado Plateau, USA: paleogeographic implications. *Sed. Geol.*, **163**, 29–66.
- Dickinson, W.R., Stair, K.N., Gehrels, G.E., Peters, L., Kowallis, B.J., Blakey, R.C., Amar, J.R.** and **Greenhalgh, B.W.** (2010) U-Pb and ⁴⁰Ar/³⁹Ar Ages for a Tephra Lens in the Middle Jurassic Page Sandstone: first Direct Isotopic Dating of a Mesozoic Eolianite on the Colorado Plateau. *J. Geol.*, **118**, 215–221.
- Do Campo, M., del Papa, C., Nieto, F., Hongn, F.** and **Petrinovic, I.** (2010) Integrated analysis for constraining palaeoclimatic and volcanic influences on clay–mineral assemblages in orogenic basins (Palaeogene Andean foreland, Northwestern Argentina). *Sed. Geol.*, **228**, 98–112.
- Doe, T.W.** and **Dott, R.H., Jr** (1980) Genetic significance of deformed cross bedding; with examples from the Navajo and Weber sandstones of Utah. *J. Sed. Petrol.*, **50**, 793–812.
- Dodd, C.D.** (1986). Sedimentology of Fluvio-Aeolian interactions, with geological examples from the British Isles. PhD Thesis, Plymouth Polytechnic.
- Dott, R.H., Byers, C.W., Fielder, G.W., Stenzel, S.R.** and **Winfrey, K.E.** (1986) Aeolian to marine transition in Cambro-Ordovician cratonic sheet sandstones of the northern Mississippi valley, U.S.A. *Sedimentology*, **33**, 345–367.
- Driese, S.G.** (1985) Interdune pond carbonates, Weber Sandstone, northern Utah and Colorado. *J. Sed. Petrol.*, **55**, 187–195.
- Dylan, B.** (1963) Blowin' in the wind. In: *The Freewheelin' Bob Dylan* (Ed. B. Dylan). Sony Music Entertainment, New York.
- Eastwood, E.N., Kocurek, G., Mohrig, D.** and **Swanson, T.** (2012) Methodology for reconstructing wind direction, wind speed and duration of wind events from aeolian cross-strata. *J. Geophys. Res.*, **117**, F03035.
- El-Fawal, F.M., El-Asmar, H.M.** and **Sarhan, M.A.E.-F.** (2011) Depositional evolution of the Middle-Upper Eocene rocks, Fayum area, Egypt. *Arabian J. Geosci.*, **2011**, 1–12.
- Elias, A.D., De Ros, L.F., Mizusaki, A.M.** and **Kawashita, K.** (2007) Isotopic evidence on the diagenetic evolution of coastal sabkha reservoirs from the Solimões Basin, northern Brazil. *Gondwana Res.*, **11**, 553–567.
- Eriksson, K.A.** and **Simpson, E.L.** (1998) Controls on spatial and temporal distribution of Precambrian eolianites. *Sed. Geol.*, **120**, 275–294.
- Eriksson, P.G., Simpson, E.L., Eriksson, K.A., Bumby, A.J., Steyn, G.L.** and **Sarkar, S.** (2000) Muddy roll-up structures in siliciclastic interdune beds of the ca. 1.8 Ga Waterberg Group, South Africa. *Palaio*, **15**, 177–183.
- Evans, D.A.D.** (2003) A fundamental Precambrian-Phanerozoic shift in earth's glacial style? *Tectonophysics*, **375**, 353–385.
- Fabuel-Perez, I., Redfern, J.** and **Hodgetts, D.** (2009) Sedimentology of an intra-montane rift-controlled fluvial dominated succession: the Upper Triassic Oukaimeden Sandstone Formation, Central High Atlas, Morocco. *Sed. Geol.*, **218**, 103–140.
- Fitzsimmons, K.E., Rhodes, E.J., Magee, J.W.** and **Barrows, T.T.** (2007) The timing of linear dune activity in the Strzelecki and Tirari deserts. *Quatern. Sci. Rev.*, **26**, 2598–2616.
- Fornós, J., Clemmensen, L.B., Gómez-Pujol, L.** and **Murray, A.S.** (2009) Late Pleistocene carbonate aeolianites on Mallorca, Western Mediterranean: a luminescence chronology. *Quatern. Sci. Rev.*, **28**, 2697–2709.
- Franceschini, G.** and **Compton, J.S.** (2004) Aeolian and marine deposits of the Tabakbaai Quarry area, western Cape, South Africa. *S. Afr. J. Geol.*, **107**, 619–632.
- Frébourg, G., Hassler, C.-A., Le Guern, P.** and **Davaud, E.** (2008) Facies characteristics and diversity in carbonate eolianites. *Facies*, **54**, 175–191.
- Fryberger, S.G.** (1979) Dune forms and wind regimes. In: *A Study of Global Sand Seas* (Ed. E.D. McKee), *U.S. Geol. Surv. Prof. Pap.*, **1052**, 137–140.
- Fryberger, S.G.** (1984) The Permian Upper Minnelusa Formation, Wyoming; ancient example of an offshore-prograding eolian sand sea with geomorphic facies, and system-boundary traps for petroleum. *Wyoming Geol. Assoc. Guidebook*, **35**, 241–271.
- Fryberger, S.G.** (1993) A review of aeolian bounding surfaces, with examples from the Permian Minnelusa Formation, USA. In: *Characterization of Fluvial and Aeolian Reservoirs* (Eds C.P. North and D.J. Prosser), *Geol. Soc. London Spec. Publ.*, **73**, 167–197.
- Fujioka, T., Chappell, J., Fifield, L.K.** and **Rhodes, E.J.** (2009) Australian desert dune fields initiated with Pliocene-Pleistocene global climatic shift. *Geology*, **37**, 51–54.
- Galusha, T.** and **Blick, J.C.** (1971) Stratigraphy of the Santa Fe Group, New Mexico. *Am. Mus. Nat. Hist. Bull.*, **144**, 127.
- Garden, R., Mentiplay, M., Cook, R.** and **Sylvester, I.** (2005) Modelling fine scale heterogeneity for optimal history matching from aeolian gas reservoirs, Neptune Field, UKCS. In: *Petroleum Geology of Northwest Europe: Proceedings of the Conference* (Eds A.G. Doré and B.A. Vining), *J. Geol. Soc. London*, **6**, 695–706.
- Garzanti, E., Ando, S., Vezzoli, G.** and **Dell'era, D.** (2003) From rifted margins to foreland basins: investigating provenance and sediment dispersal across desert Arabia (Oman, UAE). *J. Sed. Res.*, **73**, 572–588.

- Garzanti, E., Andó, S., Vezzoli, G., Lustrino, M., Boni, M. and Vermeesch, P.** (2012) Petrology of the Namib Sand Sea: long-distance transport and compositional variability in the wind-displaced Orange Delta. *Earth-Sci. Rev.*, **112**, 173–189.
- Geuna, S.E., Escosteguy, L.D. and Limarino, O.** (2010) Paleomagnetism of the Carboniferous-Permian Patquía Formation, Paganzo basin, Argentina: implications for the apparent polar wander path for South America and Gondwana during the Late Palaeozoic. *Geol. Acta*, **8**, 373–397.
- Ghinassi, M., Magi, M., Sagri, M. and Singer, B.S.** (2004) Arid climate 2.5 Ma in the Plio-Pleistocene Valdarno Basin (Northern Apennines, Italy). *Palaeogeogr. Palaeoclimatol. Palaeoecol.*, **207**, 37–57.
- Gilmer, G.J.** (2008) Paleoenvironmental interpretation of the Lower Taylor Group, Olympus Range Area, Southern Victoria Land, Antarctica. MS Thesis, University of Canterbury, 201 pp.
- Glennie, K.W.** (1970) *Desert Sedimentary Environments*. *Dev. Sedimentol.*, **14**, 222.
- Glennie, K.W.** (1972) Permian Rotliegendes of Northwest Europe interpreted in the light of modern desert sedimentation studies. *AAPG Bull.*, **56**, 1048–1071.
- Glennie, K.W.** (1998) Lower Permian – Rotliegend. In: *Petroleum Geology of the North Sea: Basic Concepts and Recent Advances* (Ed. K.W. Glennie), pp. 137–173. Blackwell Science, Oxford, UK.
- Glennie, K.W.** (2005) *The Desert of Southeast Arabia*. GeoArabia, Manama, Bahrain, 215 pp.
- Glennie, K.W. and Buller, A.T.** (1983) The Permian Weissliegend of North West Europe. The partial deformation of aeolian dune sands caused by the Zechstein transgression. *Sed. Geol.*, **35**, 43–81.
- Glennie, K. and Hurst, A.** (2007) Fluidization and associated soft-sediment deformation in eolian sandstones: Hopeman Sandstone (Permian), Scotland, and Rotliegend, North Sea. In: *Sand Injectites: Implications for Hydrocarbon Exploration and Production* (Eds A. Hurst and J. Cartwright), *AAPG Mem.*, **87**, 245–252.
- Goso, C., Piñeiro, G., DeSantaAna, H., Rojas, A., Verde, M. and Alves, C.** (2001) Caracterización estratigráfica de los depósitos continentales cuspidales neopérmicos (Formaciones Yaguari y Buena Vista) en el borde oriental de la Cuenca Norte Uruguaya. In: *Congreso Latinoamericano de Geología*, **11**. Actas, Montevideo, 18 pp. (CDROM)
- Grey, K., Hocking, R.M., Stevens, M.K., Bagas, I., Carlsen, G.M., Irimies, F., Pirajno, F., Haines, P.W. and Apak, S.N.** (2005) Lithostratigraphic nomenclature of the Officer Basin and correlative parts of the Paterson Orogen, Western Australia. Western Australia Geological Survey, Report **93**, 89 pp.
- Guern, P.L. and Davaud, E.** (2005) Recognition of ancient carbonate wind deposits: lessons from a modern analogue, Chrissi Island, Crete. *Sedimentology*, **52**, 915–926.
- Haddon, I.G. and McCarthy, T.S.** (2005) The Mesozoic-Cenozoic interior sag basins of Central Africa: the Late-Cretaceous-Cenozoic Kalahari and Okavango basins. *J. Afr. Earth Sci.*, **43**, 316–333.
- Han, Q., Qu, J., Dong, Z., Zu, R., Zhang, K., Wang, H. and Xie, S.** (2014) The effect of air density on sand transport structures and the adobe abrasion profile: a field wind-tunnel experiment over a wide range of altitude. *Bound.-Lay. Meteorol.*, **150**, 299–317.
- Hanebuth, T.J.J., Mersmeyer, H., Kudrass, H.R. and Westphal, H.** (2013) Aeolian to shallow-marine shelf architecture off a major desert since the Late Pleistocene (northern Mauritania). *Geomorphology*, **203**, 132–146.
- Hartley, A.J., Weissmann, G.S., Nichols, G.J. and Warwick, G.L.** (2010) Large distributive fluvial systems: characteristics, distribution, and controls on development. *J. Sed. Res.*, **80**, 167–183.
- Havholm, K.G. and Kocurek, G.** (1994) Factors controlling aeolian sequence stratigraphy: clues from super bounding surface features in the Middle Jurassic Page Sandstone. *Sedimentology*, **41**, 913–934.
- Havholm, K.G., Blakey, R.C., Capps, M., Jones, L.S., King, D.D. and Kocurek, G.** (1993) Aeolian genetic stratigraphy: an example from the Middle Jurassic Page sandstone, Colorado Plateau. In: *Aeolian Sediments: Ancient and Modern* (Eds K. Pye and N. Lancaster), *Int. Assoc. Sedimentol. Spec. Publ.*, **16**, 87–108.
- Hearty, P.J., Miller, G.H., Stearns, C.E. and Szabo, B.J.** (1986) Aminostratigraphy of Quaternary shorelines in the Mediterranean basin. *Geol. Soc. Am. Bull.*, **97**, 850–858.
- Hernández, R. and Echevarría, L.** (2009) Faja plegada y corrida subandina del noroeste argentino: estratigrafía, geometría y cronología de la deformación. *Rev. Asoc. Geol. Argentina*, **65**, 68–80.
- Herries, R.D.** (1993) Contrasting styles of fluvio-aeolian interaction at a downwind erg margin: Jurassic Kayenta-Navajo transition, northern Arizona, USA. In: *Characterization of Fluvial and Aeolian Reservoirs* (Eds C.P. North and D.J. Prosser), *Geol. Soc. London Spec. Publ.*, **73**, 199–218.
- Hesp, P.A.** (1999) The beach backshore and beyond. In: *Handbook of Beach and Shoreface Morphodynamics* (Ed. A.D. Short), pp. 145–169. John Wiley & Sons Ltd, Chichester, UK.
- Hesse, P.** (2011) Sticky dunes in a wet desert: formation, stabilisation and modification of the Australian desert dunefields. *Geomorphology*, **134**, 309–325.
- Hillaire-Marcel, C., Gariépy, C., Ghaleb, B., Goy, J.L., Zazo, C. and Cuerda, J.** (1996) U-Series measurements in Tyrrhenian deposits from Mallorca. Further evidence for two last-interglacial high sea levels in the Balearic Islands. *Quatern. Sci. Rev.*, **15**, 53–62.
- Holzförster, F.** (2007) Lithology and depositional environments of the lower Jurassic Clarens Formation in the Eastern Cape, South Africa. *S. Afr. J. Geol.*, **110**, 543–560.
- Horne, R.R.** (1971) Aeolian cross-stratification in Devonian of Dingle peninsula, County Kerry, Ireland. *Geol. Mag.*, **108**, 151–158.
- Hou, B., Alley, N.F., Frakes, L.A., Stoian, L. and Cowley, W.M.** (2006) Eocene stratigraphic succession in the Eucla Basin of South Australia and correlation to major regional sea-level events. *Sed. Geol.*, **183**, 297–319.
- Howell, J.A. and Mountney, N.P.** (1997) Climatic cyclicity and accommodation space in arid and semi-arid depositional systems: an example from the Rotliegende Group of the southern North Sea. In: *Petroleum Geology of the Southern North Sea: Future Potential* (Eds C.P. North and J.D. Prosser), *Geol. Soc. London Spec. Publ.*, **123**, 199–218.
- Howell, J.A. and Mountney, N.P.** (2001) Aeolian grain flow architecture: hard data for reservoir models and implications for red bed sequence stratigraphy. *Petrol. Geosci.*, **7**, 51–56.

- Howell, J.A., Schwarz, E., Spalletti, L.A. and Veiga, G.D.** (2005) The Neuquén Basin, Argentina: an overview. In: *The Neuquén Basin: A Case Study in Sequence Stratigraphy and Basin Dynamics* (Eds G.D. Veiga, L.A. Spalletti, J.A. Howell and E. Schwarz), *Geol. Soc. London Spec. Publ.*, **252**, 1–14.
- Hunter, R.E. and Rubin, D.M.** (1983) Interpreting cyclic cross-bedding, with an example from the Navajo Sandstone. In: *Eolian Sediments and Processes, Developments in Sedimentology 38* (Eds M.E. Brookfield and T.S. Ahlbrandt), pp. 429–454. Elsevier, Amsterdam.
- Hurst, A. and Glennie, K.W.** (2008) Mass-wasting of ancient aeolian dunes and sand fluidization during a period of global warming and inferred brief high precipitation: the Hopeman Sandstone (late Permian), Scotland. *Terra Nova*, **20**, 274–279.
- Irigoyen, M.V., Buchan, K.L. and Brown, R.L.** (2000) Magnetostratigraphy of Neogene Andean foreland-basin strata, lat 33 S, Mendoza Province, Argentina. *Geol. Soc. Am. Bull.*, **112**, 803–816.
- Isbell, J.L., Henry, L.C., Gulbranson, E.L., Limarino, C.O., Fraiser, M.L., Koch, Z.J., Ciccioli, P.L. and Dineen, A.A.** (2012) Glacial paradoxes during the late Paleozoic ice age: evaluating the equilibrium line altitude as a control on glaciation. *Gondwana Res.*, **22**, 1–19.
- Jennings, J.N.** (1975) Desert dunes and estuarine fill in the Fitzroy estuary (North-Western Australia). *Catena*, **2**, 215–262.
- Jerram, D.A., Mountney, N.P. and Stollhofen, H.** (1999a) Facies architecture of the Etjo Sandstone Formation and its interaction with the basal Etendeka Flood Basalts of Northwest Namibia; implications for offshore prospectivity. In: *The Oil and Gas Habitats of the South Atlantic* (Eds N.R. Cameron, R.H. Bate and V.S. Clure), *Geol. Soc. London Spec. Publ.*, **153**, 367–380.
- Jerram, D.A., Mountney, N.P., Holzforster, F. and Stollhofen, H.** (1999b) Internal stratigraphic relationships in the Etendeka Group in the Huab Basin, NW Namibia: understanding the onset of flood volcanism. *J. Geodyn.*, **28**, 393–418.
- Jerram, D.A., Mountney, N.P., Howell, J.A. and Stollhofen, H.** (2000a) The fossilised desert; recent developments in our understanding of the Lower Cretaceous deposits in the Huab Basin, NW Namibia. In: *Henno Martin Commemorative Volume* (Ed. R.M. Miller), *Com. Geol. Surv. S. W. Africa/Namibia*, **12**, 269–278.
- Jerram, D.A., Mountney, N.P., Howell, J.A., Long, D. and Stollhofen, H.** (2000b) Death of a sand sea: an active aeolian erg systematically buried by the Etendeka flood basalts of NW Namibia. *J. Geol. Soc. London*, **157**, 513–516.
- Jerzykiewicz, T., Currie, P.J., Eberth, D.A., Johnston, P.A., Koster, E.H. and Zheng, J.-J.** (1993) Djadokhta Formation correlative strata in Chinese Inner Mongolia: an overview of the stratigraphy, sedimentary geology, and paleontology and comparisons with the type locality in the pre-Altai Gobi. *Can. J. Earth Sci.*, **30**, 2180–2195.
- Jiang, X., Pan, Z., Xu, J., Li, X., Xie, G. and Xiao, Z.** (2008) Late Cretaceous aeolian dunes and reconstruction of palaeo-wind belts of the Xinjiang Basin, Jiangxi Province, China. *Palaeogeogr. Palaeoclimatol. Palaeoecol.*, **257**, 58–66.
- Jones, B.G.** (1972) Deformation structures in siltstone resulting from migration of an Upper Devonian aeolian dune. *J. Sed. Petrol.*, **42**, 935–940.
- Jones, L.S. and Blakey, R.C.** (1993) Erosional remnants and adjacent unconformities along an eolian-marine boundary in the Page Sandstone and Carmel Formation, Middle Jurassic, south-central Utah. *J. Sed. Petrol.*, **63**, 852–859.
- Jordan, O.D. and Mountney, N.P.** (2010) Styles of interaction between aeolian, fluvial and shallow marine environments in the Pennsylvanian to Permian lower Cutler beds, south-east Utah, USA. *Sedimentology*, **57**, 1357–1385.
- Jordan, O.D. and Mountney, N.P.** (2012) Sequence stratigraphic evolution and cyclicity of an ancient coastal desert system: the Pennsylvanian-Permian Lower Cutler Beds, Paradox Basin, Utah, U.S.A. *J. Sed. Res.*, **82**, 755–780.
- Kamola, D.L. and Huntoon, J.E.** (1994) Changes in rate of transgression across the Permian White Rim Sandstone, Southern Utah. *J. Sed. Res.*, **64**, 202–210.
- Kent, D.V. and Tauxe, L.** (2005) Corrected Late Triassic latitudes for continents adjacent to the North Atlantic. *Science*, **307**, 240–244.
- Klijn, J.A.** (1990) Dune forming factors in a geographical context. *Catena Suppl.*, **18**, 1–14.
- Kocurek, G.** (1980) Significance of bounding surfaces, interdune deposits and dune stratification types in ancient erg reconstruction. Doctoral Thesis, University of Wisconsin at Madison, 369 pp.
- Kocurek, G.** (1981a) Erg reconstruction: the Entrada Sandstone (Jurassic) of northern Utah and Colorado. *Palaeogeogr. Palaeoclimatol. Palaeoecol.*, **36**, 125–153.
- Kocurek, G.** (1981b) Significance of interdune deposits and bounding surfaces in eolian dune sands. *Sedimentology*, **28**, 753–780.
- Kocurek, G.** (1988) Late Paleozoic and Mesozoic eolian deposits of the western interior of the United States. *Sed. Geol.*, **56**, 413.
- Kocurek, G.** (1991) Interpretation of ancient eolian sand dunes. *Annu. Rev. Earth Planet. Sci.*, **19**, 43–75.
- Kocurek, G.** (1998) Aeolian system response to external forcing factors – a sequence stratigraphic view of the Saharan Region. In: *Quaternary Deserts and Climatic Change* (Eds A.S. Alsharan, K.W. Glennie, G.L. Whittle and C.G.S.C. Kendall), pp. 327–338. Balkema, Rotterdam/Brookfield.
- Kocurek, G.** (1999) The aeolian rock record (Yes, Virginia, it exists but it really is rather special to create one). In: *Aeolian Environments Sediments and Landforms* (Eds A.S. Goudie, I. Livingstone and S. Stokes), pp. 239–259. John Wiley and Sons Ltd, Chichester.
- Kocurek, G. and Dott, R.H.** (1983) Jurassic palaeogeography and paleoclimate of the central and southern Rocky Mountain region. In: *Mesozoic Paleogeography of the West-Central United States, Rocky Mountain Paleogeography Symposium 2* (Eds M.W. Reynolds and E.D. Dolly), pp. 101–118. SEPM, Denver.
- Kocurek, G. and Havholm, K.G.** (1993) Eolian sequence stratigraphy—a conceptual framework. In: *Siliciclastic Sequence Stratigraphy* (Eds P. Weimer and H. Posamentier), *AAPG Mem.*, **58**, 393–409.
- Kocurek, G. and Kirkland, B.L.** (1998) Getting to the source: Aeolian influx to the Permian Delaware basin region. *Sed. Geol.*, **117**, 143–149.
- Kocurek, G. and Lancaster, N.** (1999) Aeolian system sediment state: theory and Mojave Desert Kelso dune field example. *Sedimentology*, **46**, 505–515.
- Kocurek, G., Knight, J. and Havholm, K.** (1991) Outcrop and semi-regional three-dimensional architecture and reconstruction of a portion of the eolian Page Sandstone (Jurassic). In: *Three-Dimensional Facies Architecture* (Eds A. Miall and N. Tyler), *SEPM*, 25–43, Tulsa, OK.

- Kocurek, G., Lancaster, N., Carr, M. and Frank, A.** (1999) Tertiary Tsondab Sandstone Formation: preliminary bedform reconstruction and comparison to modern Namib Sand Sea dunes. *J. Afr. Earth Sci.*, **29**, 629–642.
- Kocurek, G., Robinson, N.I. and Sharp, J.M.J.** (2001) The response of the water table in coastal aeolian systems to changes in sea level. *Sed. Geol.*, **139**, 1–13.
- Kocurek, G., Carr, M., Ewing, R., Havholm, K.G., Nagar, Y.C. and Singhvi, A.K.** (2007) White Sands Dune Field, New Mexico: age, dune dynamics, and recent accumulations. *Sed. Geol.*, **197**, 313–331.
- Koning, D.J., Scott, B.A. and Dunbar, N.** (2004) *Middle-Upper Miocene Stratigraphy of the Velarde Graben, North-Central New Mexico: Tectonic and Paleogeographic Implications*. New Mexico Geological Society Guidebook, 55th Field Conference, Geology of the Taos Region, pp. 359–373. New Mexico Geological Society, New Mexico.
- Lancaster, N.** (1989) *The Namib Sand Sea: Dune Forms, Processes, and Sediments*. A.A. Balkema, Rotterdam, 200 pp.
- Lancaster, N.** (1999) Geomorphology of desert sand seas. In: *Aeolian Environments, Sediments and Landforms* (Eds A.S. Goudie, I. Livingstone and S. Stokes), pp. 49–70. John Wiley & Sons, Chichester.
- Lancaster, N.** (2000) Eolian deposits. In: *Cenozoic of Southern Africa* (Eds T.C. Partridge and R.M. Maud), *Oxford Monographs on Geology and Geophysics*, **40**, 73–87. Oxford University Press, New York.
- Lancaster, N.** (2008) Desert dune dynamics and development: insights from luminescence dating. *Boreas*, **37**, 559–573.
- Lancaster, N.** (2013) 11.12 Sand Seas and Dune fields. In: *Treatise on Geomorphology* (Ed. J.F. Shroder), pp. 219–245. Academic Press, San Diego.
- Lancaster, N., Kocurek, G., Singhvi, A.K., Pandey, V., Deynoux, M., Ghienne, J.-P. and Lo, K.** (2002) Late Pleistocene and Holocene dune activity and wind regimes in the western Sahara of Mauritania. *Geology*, **30**, 991–994.
- Langford, R.P. and Chan, M.A.** (1988) Flood surfaces and deflation surfaces within the Cutler Formation and Cedar Mesa Sandstone (Permian), southeastern Utah. *Geol. Soc. Am. Bull.*, **100**, 1541–1549.
- Langford, R.P. and Chan, M.A.** (1989) Fluvial–aeolian interactions: part II, ancient systems. *Sedimentology*, **36**, 1037–1051.
- Langford, R.P., Pearson, K.M., Duncan, K.A., Tatum, D.M., Adams, L. and Depret, P.A.** (2008) Eolian topography as a control on deposition incorporating lessons from modern dune seas: Permian Cedar Mesa Sandstone, SE Utah, U.S.A. *J. Sed. Res.*, **78**, 410–422.
- Leighton, C.L., Bailey, R.M. and Thomas, D.S.G.** (2013) The utility of desert sand dunes as Quaternary chronostratigraphic archives: evidence from the northeast Rub' al Khali. *Quatern. Sci. Rev.*, **78**, 303–318.
- Lindquist, S.J.** (1988) Practical characterisation of eolian reservoirs for development: Nugget Sandstone, Utah–Wyoming Thrust Belt. *Sed. Geol.*, **56**, 315–339.
- Liversage, D. and Robinson, D.** (1993) Prehistoric settlement and landscape development in the sandhill belt of southern Thy. *J. Danish Archaeol.*, **11**, 39–56.
- Livingstone, I., Bristow, C., Bryant, R.G., Bullard, J., White, K., Wiggs, G.F.S., Baas, A.C.W., Bateman, M.D. and Thomas, D.S.G.** (2010) The Namib Sand Sea digital database of aeolian dunes and key forcing variables. *Aeolian Res.*, **2**, 93–104.
- Lomax, J., Hilgers, A. and Radtke, U.** (2011) Palaeoenvironmental change recorded in the palaeodunefields of the western Murray Basin, South Australia: new data from single grain OSL-dating. *Quatern. Sci. Rev.*, **30**, 723–736.
- Loope, D.B.** (1985) Episodic deposition and preservation of eolian sands: a Late Palaeozoic example from southeastern Utah. *Geology*, **13**, 73–76.
- Loope, D.B. and Simpson, E.L.** (1992) Significance of thin sets of eolian cross-strata. *J. Sed. Petrol.*, **62**, 849–859.
- Loope, D.B. and Rowe, C.M.** (2003) Long-lived pluvial episodes during deposition of the Navajo Sandstone. *J. Geol.*, **111**, 223–232.
- Loope, D.B., Mason, J.A. and Dingus, L.** (1999) Lethal sandslides from eolian dunes. *J. Geol.*, **107**, 707–713.
- Loope, D., Eisenberg, L. and Waiss, E.** (2004) Navajo sand sea of near-equatorial Pangea: tropical westerlies, slumps, and giant stromatolites. In: *Field Trips in the Southern Rocky Mountains, USA* (Eds P. Nelson and E.A. Erslev), *Geol. Soc. Am. Field Guide*, **5**, 1–13.
- Loope, D.B., Seiler, W.M., Mason, J.A. and Chan, M.A.** (2008) Wind scour of Navajo Sandstone at the Wave (central Colorado Plateau, U.S.A.). *J. Geol.*, **116**, 173–183.
- Lovell, M.A., Jackson, P.D., Harvey, P.K. and Flint, R.C.** (2006) High-resolution petrophysical characterization of samples from an aeolian sandstone: the Permian Penrith sandstone of NW England. In: *Fluid Flow and Solute Movement in Sandstones: The Onshore UK Permo-Triassic Red Bed Sequence Barker* (Eds R.D. Barker and J.H. Tellam), *Geol. Soc. London Spec. Publ.*, **263**, 49–63.
- Lucas, S.G. and Cather, S.M.** (2003) *Stratigraphy of the Paleogene Chuska Sandstone, New Mexico-Arizona*. New Mexico Geological Society Guidebook, 54th Field Conference, Geology of the Zuni Plateau, pp. 389–395.
- Mack, G.H. and Dinterman, P.A.** (2002) Depositional environments and paleogeography of the Lower Permian (Leonardian) Yeso and correlative formations in New Mexico. *Mt. Geol.*, **39**, 75–88.
- MacNaughton, R.B., Dalrymple, R.W. and Narbonne, G.M.** (1997) Multiple orders of relative sea-level change in an earliest Cambrian passive-margin succession, Mackenzie Mountains, northwestern Canada. *J. Sed. Res.*, **67**, 622–637.
- Mader, D.** (1985) Aeolian sedimentation in the middle Buntsandstein in the Eifel North-South depression zone: summary of the variability of sedimentary processes in a Buntsandstein erg as a base for evaluation of the mutual relationships between aeolian sand seas and fluvial river systems in the Mid-European Buntsandstein. In: *Aspects of Fluvial Sedimentation in the Lower Triassic Buntsandstein of Europe* (Ed. D. Mader), *Lect. Notes Earth Sci.*, **004**, 90–126.
- Mader, D. and Laming, D.J.C.** (1985) Braidplain and alluvial fan environmental history and climatological evolution controlling origin and destruction of aeolian dune fields and governing overprinting of sand seas and river plains by calcrete pedogenesis in the Permian and Triassic of South Devon (England). In: *Aspects of Fluvial Sedimentation in the Lower Triassic Buntsandstein of Europe* (Ed. D. Mader), *Lect. Notes Earth Sci.*, **004**, 519–528.
- Mader, N.K. and Redfern, J.** (2011) A sedimentological model for the continental Upper Triassic Tadrart Oudou sandstone member: recording an interplay of climate and tectonics (Argana valley; south-west Morocco). *Sedimentology*, **58**, 1247–1282.

- Madsen, A.T., Murray, A.S. and Andersen, T.J.** (2007) Optical dating of dune ridges on Rømø, a barrier island in the Wadden Sea, Denmark. *J. Coastal Res.*, **23**, 1259–1269.
- Malhame, P.** (2007) Quartzarenites of the uppermost Cambrian lowermost Ordovician Kamouraska Formation, Québec, Canada: gravity flow deposits of eolian sand in the deep sea. Master Thesis, Department of Earth and Planetary Sciences, McGill University Montréal, Québec, 106 pp.
- Maretto, H., Carbone, O., Gazzera, C. and Schiuma, M.** (2002) Los reservorios de la Formación Tordillo. In: *Rocas Reservorio de las Cuencas Productivas Argentinas* (Eds M. Schiuma, G. Hinterwimmer and G. Vergani), pp. 335–358. Instituto Argentino del Petróleo y el Gas, Buenos Aires.
- Mankiewicz, P.J., Pottorf, R.J., Kozar, M.G. and Vrolijk, P.** (2009) Gas geochemistry of the Mobile Bay Jurassic Norphlet formation: thermal controls and implications for Reservoir connectivity. *AAPG Bull.*, **93**, 1319–1346.
- Marshall, J.E.A., Rogers, D.A. and Whiteley, M.J.** (1996) Devonian marine incursions into the Orcadian Basin. *J. Geol. Soc. London*, **153**, 451–466.
- Martin, D.Mc.B. and Thorne, A.M.** (2002) *Revised Lithostratigraphy of the Mesoproterozoic Bangemall Supergroup on the Edmund and Turee Creek 1:250 000 Map Sheets, Western Australia*. Record 2002/15, Department of Mineral and Petroleum Resources. Government of Western Australia, Perth, 27 pp.
- Master, S., Bekker, A. and Hofmann, A.** (2010) A review of the stratigraphy and geological setting of the Palaeoproterozoic Magondi Supergroup, Zimbabwe – type locality for the Lomagundi carbon isotope excursion. *Precambrian Res.*, **182**, 254–273.
- McClay, K.R., Norton, M.G., Coney, P. and Davis, G.H.** (1986) Collapse of the Caledonian orogen and the Old Red Sandstone. *Nature*, **323**, 147–149.
- McKee, E.D.** (1966) Structures of dunes at White Sands National Monument, New Mexico (and a comparison with structures of dunes from other selected areas). *Sedimentology*, **7**, 1–69.
- McKee, E.D.** (1979) A Study of Global Sand Seas. United States Department of the Interior, *U.S. Geol. Surv. Prof. Pap.*, **1052**, 429.
- Meadows, N.S.** (2006) The correlation and sequence architecture of the Ormskirk Sandstone Formation in the Triassic Sherwood Sandstone Group of the East Irish Sea Basin, NW England. *Geol. J.*, **41**, 93–122.
- Mello, M.R., Azambuja, N., Mohriak, W.U., Catto, A.J. and Francolin, J.B.** (2011) Promising giant new hydrocarbon frontier: the Namibian continental margin. *GeoExpro*, **8**, 64–69.
- Melvin, J. and Heine, C.J.** (2004) Sequence stratigraphy of an eolian gas sand: layering in the Permian Unayzah-A reservoir at south Ghawar, Eastern Saudi Arabia. *GeoArabia*, **9**, 103.
- Melvin, J., Wallick, B.P. and Heine, C.J.** (2010) Advances in Arabia stratigraphy: allostratigraphic layering related to paleo-water table fluctuations in eolian sandstones of the Permian Unayzah A reservoir, South Haradh, Saudi Arabia. *GeoArabia*, **15**, 55–86.
- Milana, J.P., Cevallos, M., Zavattieri, A., Pramparo, M. and Papu, H.** (1993) La secuencia terciaria de Pachaco: sedimentología, edad, correlaciones y significado paleogeográfico. *XII Congreso Geológico Argentino, Actas* **1**, 226–234.
- Milson, J.A., Mercadier, C.G.L., Livera, S.E. and Peters, J.M.** (1996) The Lower Paleozoic of Oman and its context in the evolution of a Gondwanan continental margin. *J. Geol. Soc. London*, **153**, 213–230.
- Morrisey, L.B., Braddy, S., Dodd, C., Higgs, K.T. and Williams, B.P.J.** (2012) Trace fossils and palaeoenvironments of the Middle Devonian Caherbla Group, Dingle Peninsula, southwest Ireland. *Geol. J.*, **47**, 1–29.
- Mountney, N.P.** (2006a) Eolian facies models. In: *Facies Models Revisited* (Eds H. Posamentier and R.G. Walker). *SEPM Mem.*, **84**, 19–83.
- Mountney, N.P.** (2006b) Periodic accumulation and destruction of aeolian erg sequences: the Cedar Mesa Sandstone, White Canyon, southern Utah. *Sedimentology*, **53**, 789–823.
- Mountney, N.P.** (2012) A stratigraphic model to account for complexity in aeolian dune and interdune successions. *Sedimentology*, **59**, 964–989.
- Mountney, N. and Howell, A.** (2000) Aeolian architecture, bedform climbing and preservation space in the Cretaceous Etjo Formation, NW Namibia. *Sedimentology*, **47**, 825–849.
- Mountney, N.P. and Thompson, D.B.** (2002) Stratigraphic evolution and preservation of aeolian dune and damp/wet interdune strata: an example from the Triassic Helsby Sandstone Formation, Cheshire Basin, UK. *Sedimentology*, **49**, 805–833.
- Mountney, N.P. and Jagger, A.** (2004) Stratigraphic evolution of an aeolian erg margin system: the Permian Cedar Mesa Sandstone, SE Utah, USA. *Sedimentology*, **51**, 713–743.
- Mountney, N.P. and Russell, A.J.** (2009) Aeolian dune-field development in a water-table controlled system: Skideararsándur, Southern Iceland. *Sedimentology*, **56**, 2107–2131.
- Mountney, N.P., Howell, J.A., Flint, S.S. and Jerram, D.A.** (1999a) Climate, sediment supply and tectonics as controls on the deposition and preservation of the aeolian-fluvial Etjo Sandstone Formation, Namibia. *J. Geol. Soc. London*, **156**, 771–779.
- Mountney, N.P., Howell, J.A., Flint, S.S. and Jerram, D.A.** (1999b) Relating eolian bounding-surface geometries to the bed forms that generated them: Etjo Formation, Cretaceous, Namibia. *Geology*, **27**, 159–162.
- Muhs, D.R.** (2004) Mineralogical maturity in dunefields of North America, Africa, and Australia. *Geomorphology*, **59**, 247–269.
- Muhs, D.R., Reynolds, R.R., Been, J. and Skipp, G.** (2003) Eolian sand transport pathways in the southwestern United States: importance of the Colorado River and local sources. *Quatern. Int.*, **104**, 3–18.
- Murray, A.S. and Clemmensen, L.B.** (2001) Luminescence dating of Holocene aeolian sand movement, Thy, Denmark. *Quatern. Sci. Rev.*, **20**, 751–754.
- Nichols, G.J.** (2005) Sedimentary evolution of the lower Clair Group, Devonian, west of Shetland; climate and sediment supply controls on fluvial, aeolian and lacustrine deposition. In: *Petroleum Geology of Northwest Europe* (Eds A.G. Doré and B.A. Vining), *Geol. Soc. London*, **6**, 957–967.
- Nielsen, K.A., Clemmensen, L.B. and Fornós, J.J.** (2004) Middle Pleistocene magnetostratigraphy and susceptibility stratigraphy: data from a carbonate Aeolian system, Mallorca, Western Mediterranean. *Quatern. Sci. Rev.*, **23**, 1733–1756.
- Nowatzki, C.H. and Kern, H.P.** (2000) The eolianites between Sanga do Cabral and Botucatu formations in Rio

- Grande do Sul State, Brazil. *An. Acad. Brasil. Cienc.*, **72**, 247–256.
- Olsen, H.** and **Larsen, P.H.** (1993) Lithostratigraphy of the continental Devonian sediments in North-East Greenland. *Bull. Grøn. Geol. Unders.*, **165**, 108.
- Øxnevad, I.E.I.** (1991) Aeolian and mixed aeolian-aqueous sedimentation in modern and ancient sub-tropical desert basins: examples from the Sahara and the Permo-Triassic of NW Europe. Doctoral Thesis, University of Bergen, Geological Institute, Department A, Norway, 2 vols, 161 pp. and 130 pp.
- Paim, P.S.G.** and **Scherer, C.M.S.** (2007) High-resolution stratigraphy and depositional model of wind- and water-laid deposits in the Ordovician Guaritas Rift (southernmost Brazil). *Sed. Geol.*, **202**, 776–795.
- Pais, J., Cunha, P.P., Pereira, D., Legoinha, P., Dias, R., Moura, D., Brum da Silveira, A., Kullberg, J.C. and González-Delgado, J.A.** (2012) *The Paleogene and Neogene of Western Iberia (Portugal), A Cenozoic record in the European Atlantic domain*. Springer, Berlin, Heidelberg, 158 pp.
- Paola, C.** and **Borgman, L.** (1991) Reconstructing random topography from preserved stratification. *Sedimentology*, **38**, 553–565.
- Pedersen, K.** and **Clemmensen, L.B.** (2005) Unveiling past aeolian landscapes: a ground-penetrating radar survey of a Holocene coastal dunefield system, Thy, Denmark. *Sed. Geol.*, **177**, 57–86.
- Pell, S.D., Chivas, A.R.** and **Williams, I.S.** (2000) The Simpson, Strzelecki and Tirari Deserts: development and sand provenance. *Sed. Geol.*, **130**, 107–130.
- Peterson, F.** (1988) Pennsylvanian to Jurassic eolian transportation systems in the western United States. *Sed. Geol.*, **56**, 207–360.
- Pickford, M.** and **Senut, B.** (1999) *Geology and Palaeobiology of the Namib Desert, Southwestern Africa*. Geological Survey of Namibia, Windhoek, Namibia, 155 pp.
- Poland, Z.A.** and **Simms, A.R.** (2012) Sedimentology of an erg to an erg-margin depositional system, the rush springs sandstone of Western Oklahoma, U.S.A.: implications for paleowinds across Northwestern Pangea during the Guadalupian (Middle Permian). *J. Sed. Res.*, **82**, 345–363.
- Porter, M.L.** (1986) Sedimentary record of erg migration. *Geology*, **14**, 497–500.
- Preusser, F., Radies, D.** and **Matter, A.** (2002) A 160,000-year record of dune development and atmospheric circulation in South Arabia. *Science*, **296**, 2018–2020.
- Pye, K.** and **Tsoar, H.** (1990) *Aeolian Sand and Sand Dunes*. Unwin Hyman Limited, London, 396 pp.
- Radies, D., Preusser, F., Matter, A.** and **Mange, M.** (2004) Eustatic and climatic controls on the development of the Wahiba Sand Sea, Sultanate of Oman. *Sedimentology*, **51**, 1359–1386.
- Rainbird, R.H.** and **Hadlari, T.** (2000). *Revised Stratigraphy and Sedimentology of the Paleoproterozoic Dubawnt Supergroup at the Northern Margin of Baker Lake basin, Nunavut Territory*. Geological Survey of Canada, Current Research 2000-C8, 9 pp.
- Rainbird, R.H., Hadlari, T., Aspler, L.B., Donaldson, J.A., LeCheminant, A.N.** and **Peterson, T.D.** (2003) Sequence stratigraphy and evolution of the Paleoproterozoic intracontinental Baker Lake and Thelon basins, western Churchill Province, Nunavut, Canada. *Precambrian Res.*, **125**, 21–53.
- Reimann, T., Tsukamoto, S., Harff, J., Osadczuk, K.** and **Frechen, M.** (2011) Reconstruction of Holocene coastal foredune progradation using luminescence dating – an example from the Swina barrier (southern Baltic Sea, NW Poland). *Geomorphology*, **132**, 1–16.
- Roberts, D.L.** (2008) Last Interglacial Hominid and Associated Vertebrate Fossil Trackways in 1049 Coastal Eolianites, South Africa. *Ichnos*, **15**, 190–207.
- Rodríguez-López, J.P.** (2008) Sedimentología y evolución del sistema desértico arenoso (erg) desarrollado en el margen occidental del Tethys durante el Cretácico Medio. Cordillera Ibérica, Provincias de Teruel y Zaragoza. PhD Thesis, *Universidad Complutense de Madrid Consejo Superior de Investigaciones Científicas (CSIC)*, 500 pp.
- Rodríguez-López, J.P., De Boer, P.L., Meléndez, N., Soria, A.R.** and **Pardo, G.** (2006) Windblown desert sands in coeval shallow marine deposits: a key for the recognition of coastal ergs in the mid-Cretaceous Iberian Basin, Spain. *Terra Nova*, **18**, 314–320.
- Rodríguez-López, J.P., Meléndez, N., De Boer, P.L.** and **Soria, A.R.** (2008) Aeolian sand sea development along the mid-Cretaceous western Tethyan margin (Spain): erg sedimentology and palaeoclimate implications. *Sedimentology*, **55**, 1253–1292.
- Rodríguez-López, J.P., Meléndez, N., de Boer, P.L.** and **Soria, A.R.** (2010) The action of wind and water in a mid-Cretaceous subtropical erg-margin system close to the Variscan Iberian Massif, Spain. *Sedimentology*, **57**, 1315–1356.
- Rodríguez-López, J.P., Meléndez, N., de Boer, P.L.** and **Soria, A.R.** (2012a) Controls on marine-erg margin cycle variability: aeolian-marine interaction in the Mid-Cretaceous Iberian Desert System, Spain. *Sedimentology*, **59**, 466–501.
- Rodríguez-López, J.P., Liesa, C.L., Van Dam, J., Lafuente, P., Arlegui, L., Ezquerro, L.** and **De Boer, P.L.** (2012b) Aeolian construction and alluvial dismantling of a fault-bounded intracontinental aeolian dune field (Teruel Basin, Spain); a continental perspective on Late Pliocene climate change and variability. *Sedimentology*, **59**, 1536–1567.
- Rodríguez-López, J.P., Meléndez, N., De Boer, P.L., Soria, A.R.** and **Liesa, C.L.** (2013) Spatial variability of multi-controlled aeolian supersurfaces in central-erg and marine-erg-margin systems. *Aeolian Res.*, **11**, 141–154.
- Romain, H.** and **Mountney, N.P.** (2014) Reconstruction of 3D eolian-dune architecture from 1D core data through application of outcrop analogue data. *AAPG Bull.*, **98**, 1–22.
- Roskin, J., Porat, N., Tsoar, H., Blumberg, D.G.** and **Zander, A.M.** (2011) Age, origin and climatic controls on vegetated linear dunes in the northwestern Negev Desert (Israel). *Quatern. Sci. Rev.*, **30**, 1649–1674.
- Rossi, G.R.** (2001) Arenisca Avilé: Facies, ambientes sedimentarios y estratigrafía de una Regresión forzada del Hauteriviense inferior de la Cuenca Neuquina. PhD Thesis, Facultad de Ciencias Naturales y Museo, Universidad Nacional de La Plata. Vol. 1: 231 pp. Vol. 2: 101 pp.
- Rubin, D.M.** (1987) Cross-bedding, bedforms and paleocurrents. In: *Concepts in Sedimentology and Palaeontology* (Ed. D.M. Rubin), *SEPM*, **1**, 187 pp.
- Rubin, D.M.** (1990) Lateral migration of linear dunes in the Strzelecki Desert, Australia. *Earth Surf. Proc. Land.*, **14**, 1–14.
- Rubin, D.M.** and **Carter, C.L.** (2006) *Bedforms and Cross-Bedding in Animation*. SEPM, Atlas Series, 2, DVD.

- Rubin, D.M. and Hunter, R.E.** (1982) Migration directions of primary and superimposed dunes inferred from compound crossbedding in the Navajo Sandstone. In: *11th International Congress on Sedimentology, Hamilton, Ontario, August 1982* (Eds J.O. Nriagu and R. Troost), *Int. Assoc. Sedimentol.*, pp. 69–70.
- Rubin, D.M. and Hunter, R.E.** (1985) Why deposits of longitudinal dunes are rarely recognized in the geologic record. *Sedimentology*, **32**, 147–157.
- Runkel, A.C., McKay, R.M. and Palmer, A.C.** (1998) Origin of a classic cratonic sheet sandstone: stratigraphy across the Sauk II-Sauk III boundary in the Upper Mississippi Valley. *Geol. Soc. Am. Bull.*, **110**, 188–210.
- Saye, S.E. and Pye, K.** (2006) Variations in chemical composition and particle size of dune sediments along the west coast of Jutland, Denmark. *Sed. Geol.*, **183**, 217–242.
- Sánchez, M.L., Rossi, J., Morra, S. and Armas, P.** (2008) Análisis Estratigráfico Secuencial de las Formaciones Huincul y Lisandro del Subgrupo Río Limay (Grupo Neuquén – Cretácico Tardío) en el Departamento El Cuy, Río Negro, Argentina. *Lat. Am. J. Sedimentol. Basin Anal.*, **15**, 1–26.
- Sarkar, S., Bose, P.K. and Eriksson, P.G.** (2011) Neoproterozoic tsunamiite: upper Bhandar Sandstone, Central India. *Sed. Geol.*, **238**, 181–190.
- Scheidt, S., Lancaster, N. and Ramsey, M.S.** (2011) Eolian dynamics and sediment mixing in the Gran Desierto, Mexico determined from thermal infrared spectroscopy and remote sensing data. *Geol. Soc. Am. Bull.*, **123**, 1628–1644.
- Scherer, C.M.S.** (2000) Eolian dunes of the Botucatu Formation (Cretaceous) in southernmost Brazil: morphology and origin. *Sed. Geol.*, **137**, 63–84.
- Scherer, C.M.S.** (2002) Preservation of aeolian genetic units by lava flows in the Lower Cretaceous of the Parana Basin, southern Brazil. *Sedimentology*, **49**, 97–116.
- Scherer, C.M.S. and Lavina, E.L.C.** (2006) Stratigraphic evolution of a fluvial-eolian succession: the example of the Upper Jurassic-Lower Cretaceous Guará and Botucatu formations, Paraná Basin, Southernmost Brazil. *Gondwana Res.*, **9**, 475–484.
- Schuster, M., Düringer, P., Ghienne, J.-F., Vignaud, P., Mackaye, H.T., Likius, A. and Brunet, M.** (2002) The age of the Sahara Desert. *Science*, **311**, 821.
- Schwarz, E., Spalletti, L.A. and Veiga, G.D.** (2011) La Formación Mulichinco (Cretácico Temprano) en la Cuenca Neuquina. In: *Geología y Recursos Naturales de la Provincia del Neuquén* (Eds H.A. Leanza, C. Arregui, E. Carbone, J.C. Danielli and J. Vallés), *Asoc. Geol. Argentina*, **12**, 131–144.
- Scotese, C.R.** (2001) *Atlas of Earth History, Volume 1, Paleogeography*. PALEOMAP Project, Arlington, Texas, 52 pp.
- Ségalen, L., Rognon, P., Pickford, M., Senut, B., Emmanuel, L., Renard, M. and Ward, J.** (2004) Reconstitution des morphologies dunaires et du régime des paléovents dans le Proto-Namib au cours du Miocène. *Bull. Soc. Géol. Fr.*, **175**, 537–546.
- Seike, K., Hasegawa, H. and Ichinnorov, N.** (2010) Preferred orientation of the trace fossil *Entradichnus meniscus* in eolian dune strata (Djadokhta Formation) at Tugrikiin Shiree, southern Mongolia and its paleoecological implications. *Island Arc*, **19**, 622–627.
- Senut, B., Pickford, M. and Mein, P.** (1995) Les falaises d'Awasi: une coupe-type pour le Cénozoïque continental de Namibie. *CR Acad. Sci. Paris*, **321**, 775–780.
- Séranne, M., Bruguier, O. and Moussavou, M.** (2008) U-Pb single zircon grain dating of present fluvial and Cenozoic aeolian sediments from Gabon: consequences on sediment provenance, reworking, and erosion processes on the Equatorial West African margin. *Bull. Soc. Géol. Fr.*, **179**, 29–40.
- Shaw, R.D., Etheridge, M.A. and Lambeck, K.** (1991) Development of the late Proterozoic to mid-Palaeozoic intracratonic Amadeus Basin in central Australia: a key to understanding tectonic forces in plate interiors. *Tectonics*, **10**, 688–721.
- Shinn, E.A.** (1973) Sedimentary accumulation along the leeward southeast coast of Qatar Peninsula, Persian Gulf. In: *The Persian Gulf* (Ed. B.H. Purser), pp. 199–209. Springer-Verlag, New York.
- Silva, B.A. and Fernandez, G.C.A.** (2004) The Piramboia and Botucatu formations (Triassic-Jurassic) in the subsurface of eastern Entre Rios Province. *Rev. Asoc. Geol. Argentina*, **59**, 141–151.
- Simpson, E.L. and Eriksson, K.A.** (1993) Thin eolianites interbedded within a fluvial and marine succession: early Proterozoic Whitworth Formation, Mount Isa Inlier, Australia. *Sed. Geol.*, **87**, 39–62.
- Simpson, E.L., Eriksson, K.A. and Muller, W.U.** (2012) 3.2 Ga eolian deposits from the Moodies Group, Barberton Greenstone Belt, South Africa: implications for the origin of first-cycle quartz sandstones. *Precambrian Res.*, **214–215**, 185–191.
- Simpson, E.L., Eriksson, K., Eriksson, P.A. and Bumby, A.J.** (2002) Eolian Dune Degradation and Generation of Massive Sandstone Bodies in the Paleoproterozoic Makgabeng Formation, Waterberg Group, South Africa. *J. Sed. Res.*, **72**, 40–45.
- Simpson, E.L. and Loope, D.B.** (1985) Amalgamated interdune deposits, White-Sands, New-Mexico. *J. Sed. Petrol.*, **55**, 361–365.
- Singhvi, A.K. and Porat, N.** (2008) Impact of luminescence dating on geomorphological and palaeoclimate research in drylands. *Boreas*, **37**, 536–558.
- Smith, R.A., Jones, N.S., Monaghan, A.A. and Arkley, S.** (2006) Fluvial and aeolian deposition in the Siluro-Devonian Swanshaw Sandstone Formation, SW Scotland. *Scot. J. Geol.*, **42**, 161–177.
- Som, S.M., Catling, D.C., Harnmeijer, J.P., Polivka, P.M. and Buick, R.** (2012) Air density 2.7 billion years ago limited to less than twice modern levels by fossil raindrop imprints. *Nature*, **484**, 359–362.
- Soria, A.R., Liesa, C.L., Rodríguez-López, J.P., Meléndez, N., de Boer, P.L. and Meléndez, A.** (2011) An Early Triassic evolving erg system (Iberian Chain, NE Spain): palaeoclimate implications. *Terra Nova*, **23**, 76–84.
- Spalletti, L.A. and Veiga, G.D.** (2007) Variability of continental depositional systems during lowstand sedimentation: an example from the Kimmeridgian of the Neuquén Basin, Argentina. *Lat. Am. J. Sedimentol. Basin Anal.*, **14**, 85–104.
- Spalletti, L.A., Limarino, C.O. and Colombo Piñol, F.** (2010) Internal anatomy of an erg sequence from the aeolian – fluvial system of the De La Cuesta Formation (Paganzo Basin, northwestern Argentina). *Geol. Acta*, **8**, 431–447.
- Spalletti, L.A., Arregui, C. and Veiga, G.D.** (2011) La Formación Tordillo y equivalentes (Jurásico Tardío) en la Cuenca Neuquina. In: *Geología y Recursos Naturales de la Provincia del Neuquén* (Eds H.A. Leanza, C. Arregui, E.

- Carbone, J.C. Danielli and J. Vallés). *Asoc. Geol. Argentina*, **12**, 99–111.
- Sprinkel, D.A., Kowallis, B.J. and Jensen, P.H.** (2011) Correlation and age of the Nugget Sandstone and Glen Canyon Group, Utah. In: *Sevier Thrust Belt: Northern and Central Utah and Adjacent Areas* (Eds D.A. Sprinkel, W.A. Yonkee and T.C. Chidsey Jr.), *Utah Geol. Assoc. Publ.*, **40**, 131–149.
- Stanesco, J.D.** (1991) Sedimentology and cyclicity in the Lower Permian De Chelly Sandstone on the Defiance Plateau; eastern Arizona. *Mt. Geol.*, **28**, 1–11.
- Steele, R.P.** (1981) Aeolian sands and sandstones. Two volumes. Unpublished PhD Thesis, University of Durham, Durham, UK.
- Stewart, W.D. and Walker, R.G.** (1980) Eolian coastal dune deposits and surrounding marine sandstones, Rocky Mountain Supergroup (Lower Pennsylvanian), southeastern British Columbia. *Can. J. Earth Sci.*, **17**, 1125–1140.
- Stokes, W.L.** (1968) Multiple parallel-truncation bedding planes – a feature of wind deposited sandstone formations. *J. Sed. Petrol.*, **38**, 510–515.
- Stokes, S. and Bray, H.E.** (2005) Late Pleistocene eolian history of the Liwa region, Arabian Peninsula. *Geol. Soc. Am. Bull.*, **117**, 1466–1480.
- Story, C.** (1998) Norphlet geology and 3-D geophysics: Fairway Field, Mobile Bay, Alabama. *Lead. Edge*, **17**, 243.
- Strömback, A., Howell, J.A. and Veiga, G.D.** (2005) The transgression of an erg – sedimentation and reworking/soft-sediment deformation of aeolian facies: the Cretaceous Troncoso Member, Neuquen Basin, Argentina. *Geol. Soc. London Spec. Publ.*, **252**, 163–183.
- Stump, T.E. and Van der Eem, J.G.** (1995) The stratigraphy, depositional environments and periods of deformation of the Wajid outcrop belt, southwestern Saudi Arabia. *J. Afr. Earth Sci.*, **21**, 421–441.
- Sun, J. and Muhs, D.R.** (2007) Mid latitude dune fields. In: *Encyclopedia of Quaternary Science* (Ed. S.A. Elias), pp. 607–626. Elsevier, Amsterdam.
- Sweet, M.L.** (1999) Interaction between aeolian, fluvial and playa environments in the Permian Upper Rotliegend Group, UK southern North Sea. *Sedimentology*, **46**, 171–187.
- Swezey, C.S.** (2006) Revisiting the age of the Sahara Desert. *Science*, **312**, 1138–1139.
- Swezey, C.S.** (2009) Cenozoic stratigraphy of the Sahara, Northern Africa. *J. Afr. Earth Sci.*, **53**, 89–121.
- Swezey, C.S., Havholm, K.G., Krezoski, G.M. and Marzolf, J.E.** (2012) Mississippian (Chesterian) eolian strata in the eastern United States. Southeastern Section. 61st Annual Meeting. *Geol. Soc. Am. Ab. Progr.*, **44**, 13.
- Talbot, M.R.** (1985) Major bounding surfaces in aeolian sandstones: a climatic model. *Sedimentology*, **32**, 257–266.
- Tamura, T., Bateman, M.D., Kodama, Y., Saitoh, Y., Watanabe, K., Yamagushi, N. and Matsumoto, D.** (2011) Building of shore-oblique transverse dune ridges revealed by ground-penetrating radar and optical dating over the last 500 years on the Tottori coast, Japan Sea. *Geomorphology*, **132**, 153–166.
- Tanner, L.H. and Lucas, S.G.** (2007) The Moenave Formation: sedimentologic and stratigraphic context of the Triassic-Jurassic boundary in the Four Corners area, southwestern U.S.A. *Palaeogeogr. Palaeoclimatol. Palaeoecol.*, **244**, 111–125.
- Tawadros, E.E.** (2012) *Geology of North Africa*. CRC Press/Balkema, Taylor & Francis Group, London, UK, 952 pp.
- Taylor, T., Stancliffe, R., Macaulay, C. and Hathon, L.** (2004) High temperature quartz cementation and the timing of hydrocarbon accumulation in the Jurassic Norphlet Sandstone, offshore Gulf of Mexico, USA. *Geol. Soc. Spec. Publ.*, **237**, 257–278.
- Tedford, R.H., Swinehart, J.B., Swisher, C.C., III, Prothero, D.R., King, S.A. and Tierney, T.E.** (2005) The Whitneyan-Arikarean transition in the High Plains. In: *The Terrestrial Eocene-Oligocene Transition in North America* (Eds D.R. Prothero and R.J. Emry), pp. 312–334. Cambridge University Press, Cambridge.
- Telfer, M.W., Bailey, R.M., Burrough, S.L., Stone, A.E.S., Thomas, D.S.G. and Wiggs, G.S.F.** (2010) Understanding linear dune chronologies: insights from a simple accumulation model. *Geomorphology*, **120**, 195–208.
- Telfer, M.W.** (2011) Growth by extension, and reworking, of a south-western Kalahari linear dune. *Earth Surf. Proc. Land.*, **36**, 1125–1135.
- Teller, J.T., Glennie, K.W., Lancaster, N. and Singhvi, A.K.** (2000) Calcareous dunes of the United Arab Emirates and Noah's Flood: the postglacial reflooding of the Persian (Arabian) Gulf. *Quatern. Int.*, **68–71**, 297–308.
- Telles Antunes, M., Elderfield, H., Legoinha, P., Nascimento, A. and Pais, J.** (1999) A stratigraphic framework for the Miocene from the lower Tagus Basin (Lisbon, Setúbal Peninsula, Portugal). Depositional sequences, biostratigraphy and isotopic ages. *Rev. Soc. Geol. Esp.*, **12**, 3–15.
- Tesón, E., Pueyo, E.L., Teixell, A., Barnolas, A., Agustí, J. and Furió, M.** (2010) Magnetostratigraphy of the Ouarzazate Basin: implications for the timing of deformation and mountain building in the High Atlas Mountains of Morocco. *Geodin. Acta*, **23**, 151–165.
- Tewes, D.W. and Loope, D.B.** (1992) Palaeo-yardangs: wind-scoured desert landforms at the Permo-Triassic unconformity. *Sedimentology*, **39**, 251–261.
- Thiry, M., Bertrand-Ayrault, M. and Grisoni, J.C.** (1988) Ground water silicification and leaching in sands: example of the Fontainebleau Sand (Oligocene) in the Paris Basin. *Geol. Soc. Am. Bull.*, **100**, 1283–1290.
- Thomas, D.S.G.** (2013) Reconstructing paleoenvironments and palaeoclimates in drylands: what can landform analysis contribute? *Earth Surf. Proc. Land.*, **38**, 3–16.
- Thomas, D.S.G. and Shaw, P.A.** (1990) The deposition and development of the Kalahari Group sediments, central southern Africa. *J. Afr. Earth Sci.*, **10**, 187–197.
- Thornton, D.A.** (1990) *A Potential Aeolian Oil Reservoir: The Mid Devonian Tandalgoo Formation, Canning Basin; Depositional Models and Diagenesis*. Degree of Master of Science, University of Adelaide.
- Tietze, K.W., Becker, M., Dietrich, B., Hoepfner, R., Karle, M., Lampe, C., Lemmer, F., Radies, D., Rose, E., Steih-Winkler, A. and Weston, N.** (1997) Ein Buntsandstein-Profil am Weststrand der Hessischen Senke (Raum Marburg) (A New Red Sandstone profile at the west edge of the Hessischen Senke, Marburg area). *Geol. Palaeontol.*, **31**, 285–294.
- Tirsgaard, H. and Øxnevad, I.E.I.** (1998) Preservation of pre-vegetational mixed fluvio-aeolian deposits in a humid climatic setting: an example from the Middle Proterozoic Eriksfjord Formation, Southwest Greenland. *Sed. Geol.*, **120**, 295–317.
- Trewin, N.H. and Fallick, A.E.** (2000) Quartz cement origins and budget in the Tumblagooda Sandstone, Western Australia. In: *Quartz Cementation in Sandstones* (Eds R.H.

- Worden and S. Morad), *Int. Assoc. Sedimentol. Spec. Publ.*, **29**, 219–229.
- Trewin, N.H. and Thirlwall, M.F.** (2002) The old red sandstone. In: *The Geology of Scotland* (Ed. N.H. Trewin), *J. Geol. Soc. London*, 213–250. London.
- Tripaldi, A. and Limarín, C.O.** (2005) Vallecito Formation (Miocene): the evolution of an eolian system in an Andean foreland basin (northwestern Argentina). *J. S. Am. Earth Sci.*, **19**, 343–357.
- Turner, P., Burley, S.D., Rey, D. and Prosser, J.** (1995) Burial history of the Penrith Sandstone (Lower Permian) deduced from the combined study of fluid inclusion and palaeomagnetic data. In: *Palaeomagnetic Applications in Hydrocarbon Exploration and Production* (Eds P. Turner and A. Turner), *Geol. Soc. London Spec. Publ.*, **98**, 43–78.
- Tyrrell, S., Leleu, S., Kate Souders, A., Haughton, P.D.W. and Stephen, D.J.** (2009) K-feldspar sand-grain provenance in the Triassic, west of Shetland: distinguishing first-cycle and recycled sediment sources? *J. Geol.*, **44**, 692–710.
- Uba, C.E., Heubeck, C. and Hulka, C.** (2005) Facies analysis and basin architecture of the Neogene Subandean synorogenic wedge, southern Bolivia. *Sed. Geol.*, **180**, 91–123.
- Umhoefer, P.J., Dorsey, R.J., Willsey, S., Mayer, L. and Renne, P.** (2001) Stratigraphy and geochronology of the Comodú Group near Loreto, Baja California sur, Mexico. *Sed. Geol.*, **144**, 125–147.
- Veiga, G. and Spalletti, L.A.** (2007) The Upper Jurassic (Kimmeridgian) fluvial-aeolian systems of the southern Neuquen Basin, Argentina. *Gondwana Res.*, **11**, 286–302.
- Veiga, G.D., Spalletti, L.A. and Flint, S.** (2002) Aeolian/fluvial interactions and high-resolution sequence stratigraphy of a non-marine lowstand wedge: the Avile Member of the Agrio Formation (Lower Cretaceous), central Neuquen Basin, Argentina. *Sedimentology*, **49**, 1001–1019.
- Veiga, G.D., Howell, J.A. and Strömbäck, A.** (2005) Anatomy of a mixed marine/non-marine lowstand wedge in a ramp setting. The record of a Barremian/Aptian complex relative sea-level fall in Central Neuquén Basin, Argentina. In: *The Neuquén Basin: A Case Study in Sequence Stratigraphy and Basin Dynamics* (Eds G.D. Veiga, L.A. Spalletti, J.A. Howell and E. Schwarz), *Geol. Soc. London Spec. Publ.*, **252**, 139–162.
- Veiga, G.D., Schwarz, E. and Spalletti, L.A.** (2011a) Análisis estratigráfico de la Formación Lotena (Calloviano superior – Oxfordiano inferior) en la Cuenca Neuquina Central, República Argentina. Integración de información de afloramientos y subsuelo. *Andean Geol.*, **38**, 171–197.
- Veiga, G.D., Schwarz, E. and Spalletti, L.A.** (2011b) El Miembro Avilé de la Formación Agrio (Cretácico Temprano). In: *Geología y Recursos Naturales de la Provincia del Neuquén* (Eds H.A. Leanza, C. Arregui, E. Carbone, J.C. Danieli and J. Vallés), *Asoc. Geol. Argentina*, **12**, 161–173.
- Vermeesch, P., Fenton, C.R., Kober, F., Wiggs, G.F.S., Bristow, C.S. and Xu, S.** (2010) Sand residence times of one million years in the Namib Sand Sea from cosmogenic nuclides. *Nat. Geosci.*, **3**, 862–865.
- Vignaud, P., Düringer, P., Mackaye, H.T., Likius, A., Blondel, C., Boisserie, J.-R., de Bonis, L., Eisenmann, V., Etienne, M.-E., Geraads, D., Guy, F., Lehmann, T., Lihoreau, F., Lopez-Martinez, N., Mourer-Chauviré, C., Otero, O., Rage, J.-C., Schuster, M., Viriot, L., Zazzo, A. and Brunet, M.** (2002) Geology and palaeontology of the Upper Miocene Toros-Menalla hominid locality, Chad. *Nature*, **418**, 152–155.
- Voss, R.** (2002) Cenozoic stratigraphy of the southern Salar de Antofalla region, northwestern Argentina. *Rev. Geol. Chile*, **29**, 167–189.
- Waichel, B.L., Scherer, C.M.S. and Frank, H.T.** (2008) Basaltic lava flows covering active aeolian dunes in the Paraná Basin in southern Brazil: features and emplacement aspects. *J. Volcanol. Geoth. Res.*, **171**, 59–72.
- Wakefield, O.J.W. and Mountney, N.P.** (2013) Stratigraphic architecture of back-filled incised-valley systems: Pennsylvanian-Permian lower Cutler beds, Utah, USA. *Sediment. Geol.*, **298**, 1–16.
- Ward, J.D.** (1988) Eolian fluvial and pan (playa) facies of the Tertiary Tsonab sandstone formation in the Central Namib Desert, Namibia. *Sed. Geol.*, **55**, 143–162.
- Wasson, R.J. and Hyde, R.** (1983) Factors determining desert dune type. *Nature*, **304**, 337–339.
- Weissmann, G.S., Hartley, A.J., Nichols, G.J., Scuderi, L.A., Olsen, M., Buehler, H. and Banteah, R.** (2010) Fluvial form in modern continental sedimentary basins: distributive fluvial systems. *Geology*, **38**, 39–42.
- Whybrow, P.J., Friend, P.F., Ditchfield, P.W. and Bristow, C.S.** (1999) Local stratigraphy of the Neogene outcrops of the coastal area: Western Region, Emirate of Abu Dhabi, United Arab Emirates. In: *Fossil Vertebrates of Arabia: With Emphasis on the Late Miocene Faunas, Geology, and Palaeoenvironments of the Emirate of Abu Dhabi, United Arab Emirates* (Eds P.J. Whybrow and A. Hill), pp. 28–37. Yale University Press, New Haven, CT.
- Williams, G.E.** (1998) Late Neoproterozoic periglacial aeolian sand sheet, Stuart Shelf, South Australia. *Aust. J. Earth Sci.*, **45**, 733–741.
- Williams, M.R.** (2009) Stratigraphy of Upper Pennsylvanian cyclic carbonate and siliciclastic rocks, western Paradox Basin, Utah. In: *The Paradox Basin Revisited – New Developments in Petroleum Systems and Basin Analysis* (Eds W.S. Houston, L.L. Wray and P.G. Moreland), pp. 381–435. Rocky Mountain Association of Geologists, Denver, CO.
- Wilson, I.G.** (1973) Ergs. *Sed. Geol.*, **10**, 77–106.
- Xie, Y., Wang, J., Jiang, X., Li, M., Xie, Z., Luo, J., Hou, G., Liu, F., Wang, Y., Zhang, M., Zhu, H., Wang, D., Sun, Y. and Cao, J.** (2005) Sedimentary characteristics of the Cretaceous desert facies in the Ordos Basin and their hydrogeological significance. *Acta Sedimentol. Sinica*, **23**, 73–83.
- Zavala, C. and Freije, H.** (2001) On the understanding of aeolian sequence stratigraphy: an example from Miocene – Pliocene deposits in Patagonia, Argentina. *Riv. Ital. Paleontol. Stratigr.*, **107**, 251–264.
- Zavala, C., Maretto, H. and Di Meglio, M.** (2005a) Hierarchy of bounding surfaces in aeolian sandstones of the Jurassic Tordillo Formation (Neuquen Basin, Argentina). *Geol. Acta*, **3**, 133–145.
- Zavala, C., Mosquera, A. and Joo Kim, H.** (2005b) *Los depósitos eólicos de la Formación Mulichinco (Valanginiano) en el área de Fortín de Piedra-Cuenca Neuquina*. VI Congreso de Exploración y Desarrollo de Hidrocarburos. Technical Papers on CD. Mar del Plata, Argentina.
- Zheng, H., Powell, Ch.Mc., Butcher, K. and Cao, J.** (2003) Late Neogene loess deposits in southern Tarim Basin:

tectonic and palaeoenvironmental implications. *Tectonophysics*, **375**, 49–59.

Zhou, L., Williams, M.A.J. and Peterson, J.A. (1994) Late Quaternary aeolianites, palaeosols and depositional environments of the Nepean Peninsula, Victoria, Australia. *Quatern. Sci. Rev.*, **13**, 225–239.

Manuscript received 29 December 2012; revision accepted 5 March 2014

Supporting Information

Additional Supporting Information may be found in the online version of this article:

Table S1. Compilation of main Archean, Palaeoproterozoic, Mesoproterozoic and Neoproterozoic aeolian sand systems.

Table S2. Compilation of main Cambrian, Ordovician, Silurian, Devonian, Carboniferous and Permian aeolian sand systems.

Table S3. Compilation of main Triassic, Jurassic and Cretaceous aeolian sand systems.

Table S4. Compilation of main Palaeogene and Neogene aeolian sand systems.

Table S5. Compilation of main Quaternary inland dunefield (ergs).

Table S6. Compilation of main Quaternary coastal dunefields.

Data S1. References text and tables.