


Ancient DNA identification of domestic animals used for leather objects in Central Asia during the Bronze Age

The Holocene
1–8
© The Author(s) 2016
Reprints and permissions:
sagepub.co.uk/journalsPermissions.nav
DOI: 10.1177/0959683616641741
hol.sagepub.com


Oskar Schröder,^{1,2} Mayke Wagner,² Saskia Wutke,¹ Yong Zhang,³
Yingxia Ma,⁴ Dongliang Xu,³ Tomasz Goslar,^{5,6} Reinder Neef,⁷
Pavel E Tarasov⁸ and Arne Ludwig¹

Abstract

The arid climate of many regions within Central Asia often leads to excellent archaeological preservation, especially in sealed funerary contexts, allowing for ancient DNA analyses. While geneticists have looked at human remains, clothes, tools, and other burial objects are often neglected. In this paper, we present the results of an ancient DNA study on Bronze Age leather objects excavated from tombs of the Wupu cemetery in the Hami Oasis and Yanghai cemetery in the Turpan Oasis, both in Xinjiang Uyghur Autonomous Region of northwestern China. In addition to species identification of goat (*Capra aegagrus/hircus*), sheep (*Ovis orientalis/aries*), and cattle (*Bos primigenius/taurus*), mitochondrial haplogroups were determined for several samples. Our results show that Bronze Age domesticated goats and sheep from the Hami and Turpan oases possessed identical or closely related haplotypes to modern domestic animals of this area. The absence of leather produced from wild animals emphasizes the importance of animal husbandry in the cultures of Wupu and Yanghai.

Keywords

ancient DNA, goat, leather, sheep, Wupu, Yanghai

Received 11 December 2015; revised manuscript accepted 25 February 2016

Introduction

Sheep, goat, and cattle were domesticated in the Fertile Crescent of southwest Asia at around 10,000 years ago (Tapio et al., 2006; Zeder, 2008), from where they spread across Eurasia, entering northwestern China about 6000–6800 years ago (Lv et al., 2015). Note that calendar ages are consistently used in this study, unless stated otherwise.

The 1st millennium BC graves of Wupu and Yanghai in the Hami and Turpan oases, respectively, of the Xinjiang Uyghur Autonomous Region have yielded well-preserved human mummies as well as tools, clothing, and other mortuary offerings (Beck et al., 2014; Guojia Wenwuju, 2012; Xinjiang tu lu fan xue yan jiu yuan and Xinjiang wen wu kao gu yan jiu suo, 2011). Their leather coats, bracers, and quivers provide a valuable tool for identifying the domesticated and wild animals, which were in human use during this time. Although some of the leather samples have been identified as sheep-derived before DNA studies, due to the presence of wool, other samples were heavily processed preventing a reliable wool/hair identification. Since the extremely arid taphonomic conditions of Xinjiang are favorable not only for leather but also for DNA preservation (Li et al., 2010, 2011), determining the species of origin using mitochondrial DNA genotyping is a promising approach.

Previously, DNA was extracted successfully from ancient leather and parchment samples in only a very few cases (Hofreiter et al., 2015; Poulakakis et al., 2007; Schlumbaum et al., 2010; Teasdale et al., 2015; Vuissoz et al., 2007). Tanning and dyeing

during the processing of hides often heavily degrade DNA (Brandt et al., 2011; Vuissoz et al., 2007) or may even introduce contaminations using animal brains as a tanning agent (Schlumbaum et al., 2010). In this study, we used sequences of two mitochondrial genes *cytochrome b* and *control region* to determine the species of origin and their mitochondrial haplogroup, addressing the biogeographic origin of animals.

¹Department of Evolutionary Genetics, Leibniz Institute for Zoo and Wildlife Research, Germany

²Eurasia Department, German Archaeological Institute, Germany

³Academia Turfanica, P.R. China

⁴Hami Museum, P.R. China

⁵Faculty of Physics, Adam Mickiewicz University in Poznań, Poland

⁶Poznan Radiocarbon Laboratory, Foundation of the A. Mickiewicz University, Poland

⁷Department of Natural Sciences, German Archaeological Institute, Germany

⁸Section Paleontology, Institute of Geological Sciences, Free University of Berlin, Germany

Corresponding author:

Oskar Schröder, Department of Evolutionary Genetics, Leibniz Institute for Zoo and Wildlife Research, Alfred-Kowalke-Str. 17, 10315 Berlin, Germany.

Email: oschroeder@izw-berlin.de

Materials and methods

Sites and samples

The Yanghai cemetery (Figure 1) is an archaeological site located in a present-day gravel desert (Jiang et al., 2006; Xinjiang tu lu fan xue yan jiu yuan and Xinjiang wen wu kao gu yan jiu suo, 2011) in the northeastern part of the Turpan (also known as Turfan) Oasis, c. 43 km southeast of the city of Turpan. The climate of this area is extremely continental, with cold dry winters, hot dry summers, and seasonal temperature amplitude of above 70°C (Domrös and Peng, 1988). Annual precipitation is below 50 mm, everywhere outside the mountain ranges. The continual aridity ensures very good preservation of all types of organic materials, even over several millennia (e.g. Beck et al., 2014; Wagner et al., 2009). Archaeological excavation conducted at the Yanghai cemetery has shown that it is relatively large (about 54,000 m²) and was used for a long period, starting in c. 12th century BC until c. 2nd century AD (Kramell et al., 2014; Xinjiang tu lu fan xue yan jiu yuan and Xinjiang wen wu kao gu yan jiu suo, 2011). Excavations of the individual tombs have shed light on the subsistence strategy of the people buried there. However, to date only a handful of these data have been published (Beck et al., 2014; Jiang et al., 2006, 2007, 2009; Kramell et al., 2014).

The Wupu cemetery (Figure 1) is known mainly for its well-preserved mummies (Wang, 1999). The site is named after Wupu village, located about 60 km west of Hami, an important oasis on the ancient Silk Road, northeast of the Tarim Basin. The dry continental environments around Wupu are similar to that of Yanghai.



Figure 1. Map showing the main physiographic features of the Xinjiang Uyghur Autonomous Region in northwestern China and the location of the Yanghai and Wupu archaeological sites northeast of the Tarim Basin.

Table 1. Leather sample data and sequencing results (cf. Figure 2).

Collection number	Locality	Archaeological object	Sample ID	Animal species	Haplogroup	GenBank accession numbers	Best match (sequence identity)
86HWM:1-00751	Wupu	Leather coat	130414/ 24	<i>Ovis orientalis</i>	Sheep A	KU686163,70,76,83,87	KR610959 (99%)
			130414/ 26	<i>Ovis orientalis</i>	Sheep intermediate A/B	KU686164,71,77,84,88	JX235856 (100%)
			130414/ 28	<i>Ovis orientalis</i>	Sheep B	KU686165,72,78,85,89	KR011777 (100%)
86HWM:2-00752	Wupu	Leather coat	130414/ 33	<i>Capra aegagrus</i>	Goat A2	KU686166,73,79,90	KP677511 (100%)
			130414/ 34	<i>Capra aegagrus</i>	Goat A3	KU686167,74,80,91	KP120681 (100%)
2003SAYIM195:9	Yanghai	Bow and arrow sheath	130417/ 82	<i>Capra aegagrus</i>	PCR fail	KU68616781	KP662716 (98%)
			130417/ 83	PCR fail	PCR fail		
			130417/ 84	<i>Bos primigenius</i>	PCR fail	KU686168,75	KT260196 (98%)
2003SAYIM195:10	Yanghai	Bow and arrow sheath	130417/ 85	<i>Ovis orientalis</i>	PCR fail	KU686169	KT781689 (100%)
2003SAYIM195:12	Yanghai	Leather bracer	130417/ 87	<i>Ovis orientalis</i>	Sheep A1a (not repeatable)	KU686182,86	KF302440 (100%)

Bolded numbers in Sample ID are used throughout the text.

The graveyard, consisting of 114 graves, was excavated during several campaigns (the last in 1991) by the archaeological team from the Hami Museum and revealed numerous mummies and artifacts assigned to the Bronze Age (c. 15th–8th century BC). Since a comprehensive excavation report has not yet been published, objects hosted in the museum partly lack a clear burial context and tomb identification.

Samples used for the DNA analysis were provided by Chinese cooperation partners representing the Chinese Academy of Cultural Heritage, Xinjiang Uygur Autonomous Region Bureau of Cultural Heritage, Xinjiang Institute of Archaeology, the Bureau of Cultural Relics of the Turpan Prefecture, and the archaeological museums in Hami and Turpan, which host archaeological objects from Wupu and Yanghai, respectively. In April 2013, leather artifacts were sampled for this study, within the framework of the Chinese–German Silk Road Fashion research project. In total, 10 selected samples (Table 1) represent one or two tombs from Wupu, excavated in 1986, and one tomb (M195) from Yanghai, excavated in 2003. The Wupu objects include two leather coats, sewn together from several smaller pieces of leather that were sampled separately with wool/hair still attached to the inner surface (Figure 2a and b). Whether they were found in one or two tombs in the Wupu cemetery could not be established based on the excavation records. The Yanghai objects consist of two leather sheaths for a bow and arrows (Figure 2c and d) and a short leather bracer (Figure 2e). Short-living herbaceous plant remains (Table 2), associated with the leather fur coat number 86HWM:2-00752 from Wupu, were radiocarbon dated to 2515 ± 30 (Poz-57373), 2480 ± 30 (Poz-57371), 2470 ± 30 (Poz-57374), and 2465 ± 30 (Poz-57375) ¹⁴C years before present (BP). Being converted to calendar years using the OxCal software (Bronk Ramsey, 2013) and IntCal13 atmospheric curve (Reimer et al., 2013), these dates translate to 762–541 BC (95.4% probability range of the four combined dates). The thread, with which the leather patches of coat number 86HWM:1-00751 were sewn together, was radiocarbon dated to 2500 ± 30 (Poz-72755) ¹⁴C yr BP (i.e. 788–537 BC, 95.4% probability). One of the leather samples analyzed in this study from tomb Yanghai M195 was radiocarbon dated to 2475 ± 30 (Poz-76927) ¹⁴C yr BP. Calibrated to calendar years, this date represents the interval 771–431 BC (95.4% probability range). Our results suggest that the leather fur coats number 86HWM:2-00752 and 86HWM:1-00751 from Wupu and the tomb M195 objects from Yanghai represent broadly the same time interval at c. 8th to 5th century BC (95.4% probability range). The calibrated radiocarbon ages of the analyzed objects from Wupu are younger than the ages previously suggested by the excavators based on typological associations. However, three dates obtained on wood samples from other tombs (Table 2) demonstrate that the Wupu graveyard was already in use during much earlier times (i.e. 1409–806 BC) in




Figure 2. The five archaeological objects examined in this study (cf. Table 1): (a) 86HWM:1-00751; (b) 86HWM:2-00752; (c) 2003SAYIM195:9 (arrows: original location of the three samples taken); (d) 2003SAYIM195:10; (e) 2003SAYIM195:12 (scale bar in centimeters, all images by D Hosner, German Archaeological Institute).

line with the dates published earlier (Wang, 1999 and references therein). The ‘old wood effect’ (see Dong et al., 2014 for discussion of this problematic in northwestern China) cannot be significant as all our sampled trees were relatively young at their death. The horse bridle made of wood and leather from the 86HWM3 tomb of Wupu revealed an age between 1108 and 901 BC (95.4% probability). This date indicates the presence of domesticated horses and horse riding in the Hami Oasis *c.* 3000 years ago and corroborates the other evidence for early horse riding from the Turpan Oasis (Beck et al., 2014; Kramell et al., 2014) and the Kunlun Mountains (Wagner et al., 2011).

Hair of seven modern sheep and three goats were also sampled for comparison from two family farmsteads in Ya’erguolecun, Turpan, Xinjiang Uyghur Autonomous Region. The sheep of both farms belonged to an indeterminate breed of brown to black fat-tailed sheep used for felt production, some of which were horned. The goats belonged to a white, horned indeterminate dairy breed. The animals were almost certainly closely related.

DNA extraction and amplification

For ancient samples, all of the following procedures up to the first PCR set-up were performed in a dedicated ancient DNA lab, while PCR and post-PCR steps were performed in a separate laboratory, according to the recommendations of Cooper and Poinar (2000). Skin and leather samples were cut to pieces of approximately 3 mm by 3 mm depending on the shape of the original sample. DNA was extracted from these pieces according to established protocol (Dabney et al., 2013). Of each sample, at least two extractions were made with an additional mock extraction per three extractions. The species of each sample was first determined by amplifying three fragments (107, 106, and 91 bp, respectively, including primers) located in the mitochondrial *cytochrome b* gene (cytB) in a two-step multiplex PCR. The first PCR included all cytB primers (Table 3) and was conducted at a 10- μ L reaction volume containing 0.5 U AmpliTaq Gold DNA Polymerase

(Thermo Fisher), 1 \times Buffer II (Thermo Fisher), 4 mM MgCl₂, 0.25 mM dNTP, 1 mg/mL BSA, 0.075 μ M of each primer, and 2 μ L DNA extract. The resulting product was then re-amplified in a second PCR including only the two primers for a single fragment at a reaction volume of 20 μ L at the same concentrations except using 0.25 U AmpliTaq Gold DNA Polymerase, 0.75 μ M of each primer, and 0.5 μ L product of the previous PCR. Both PCRs were run at initial denaturation of 94°C for 10 min, at 30 cycles with 94°C for 30 s, 57°C for 45 s, and 72°C for 45 s, and a final extension at 72°C for 5 min. The product of the second PCR was checked on a 3% agarose gel and sequenced on a 3130XL Genetic Analyzer (Applied Biosystems). After genera identification, specific primers (Table 3) were employed to amplify fragments of the mitochondrial control region (*Ovis* (sheep): three fragments 144, 140, and 140 bp in length, *Capra* (goat) 170 bp, *Bos* (cattle) 177 bp including primers) using the same concentrations and programs as before, except for sample 130417/84 where no BSA was added.

DNA from modern sheep and goat (hair samples) was extracted using the First Magnetic Forensic Kit (GEN-IAL) according to the manual. After extraction, only the second PCR step was performed for control region fragments, followed by Sanger sequencing as described before.

DNA analysis

Sequences were aligned in BioEdit. Haplogroups were identified with BLAST searches and comparison to reference sequences on DOME (Peng et al., 2015). For comparison, mitochondrial control region sequences from different geographical locations were downloaded from GenBank (sheep: KF938317–59, HM236176–77, JX235856; argali: HM236188, JX101654; urial: NC_026064, KF938361; Asiatic mouflon: NC_026063, KF938360, KF312238; goat: DQ089106–13, 16, 35, 47, 55–59, 86–88, 91, AJ317760–1, EF617863–76, GQ342248, DQ847506–09, EF167788–802; Siberian ibex: NC_020626, FJ207529) to

Table 2. AMS radiocarbon dates processed on samples from the Wupu and Yanghai tombs and discussed in this study. All dates were generated in the Poznan Radiocarbon Laboratory, Poland. The dates expressed in ^{14}C BP (radiocarbon years before AD 1950, conventionally taken as the 'present') were converted to calendar years (BC) using OxCal v4.2.4 calibration software (Bronk Ramsey, 2013) and atmospheric data from Reimer et al. (2013). Given are intervals of calendar ages, where the true ages of the samples encompass the probability ranges of c. 68% and c. 95%. Radiocarbon-dated plant material identification has been done by R Neef. One leather sample (Sample ID 130417/86) from the Yanghai tomb M195 disintegrated in chemical preparation almost completely, and extremely little amount of carbon (0.01 mg) obtained from it did not allow for a reliable date. For this reason, the reported date 3400 ± 170 ^{14}C BP (Poz-76845) is regarded as 'very poor' and is not recommended for further use.

Sample ID	Laboratory number	Cemetery and tomb number	Dated material/archaeological context	Radiocarbon date, ^{14}C BP ($\pm 1\sigma$)	Calibrated individual dates, cal. BC (68.2% probability)	Calibrated individual dates, cal. BC (95.4% probability)
130414/36	Poz-57373	86HWM:2-00752, Wupu	Barley (<i>Hordeum vulgare</i>) spikelets/leather fur coat	2515 ± 30	775 (15.4%) 747 BC 685 (10.4%) 666 BC 642 (42.4%) 555 BC	793 (26.6%) 727 BC 719 (1.8%) 705 BC 695 (66.9%) 541 BC
130414/35	Poz-57371	86HWM:2-00752, Wupu	Brome grass (<i>Bromus</i> sp.) seeds/leather fur coat	2480 ± 30	756 (11.1%) 728 BC 717 (3.6%) 706 BC 694 (5.6%) 679 BC 671 (47.9%) 542 BC	774 (94.9%) 482 BC 441 (0.5%) 434 BC
130414/37	Poz-57374	86HWM:2-00752, Wupu	Non-identified herbaceous plant stem/thorn fragment/leather fur coat	2470 ± 30	753 (27.1%) 682 BC 669 (20.8%) 612 BC 593 (20.3%) 538 BC	768 (92.4%) 476 BC 464 (1.2%) 453 BC 445 (1.8%) 431 BC
130414/38	Poz-57375	86HWM:2-00752, Wupu	Burdock (<i>Arctium</i> sp.) burr/leather fur coat	2465 ± 30	751 (26.9%) 683 BC 669 (12.4%) 636 BC 626 (3.5%) 614 BC 592 (25.4%) 516 BC	764 (95.4%) 430 BC
	Combined 4 dates	86HWM:2-00752, Wupu			752 (11.3%) 735 BC 689 (26.5%) 613 BC 592 (30.3%) 546 BC	762 (19.1%) 705 BC 695 (76.3%) 541 BC
130414/30	Poz-72755	86HWM:1-00751, Wupu	Sewing wool thread/leather fur coat	2500 ± 30	767 (11.1%) 744 BC 687 (10.8%) 665 BC 644 (46.4%) 552 BC	788 (95.4%) 537 BC
130415/55	Poz-57377	86HWM3:1, Wupu	Wood sample/wooden cheek-piece of a horse bridle	2825 ± 30	1012 (68.2%) 927 BC	1108 (0.2%) 1105 BC 1081 (1.3%) 1065 BC 1056 (93.9%) 901 BC
130415/52	Poz-57378	86HWMNN:1, Wupu	Blue wool thread/long robe	2465 ± 30	752 (25.8%) 686 BC 667 (11.6%) 636 BC 622 (2.4%) 614 BC 595 (28.3%) 515 BC	760 (27.8%) 682 BC 671 (56.7%) 480 BC 469 (10.9%) 414 BC
130416/79	Poz-57381	No tomb attribution, Wupu	Wood/poplar trunk from the tomb chamber	3050 ± 30	1385 (39.6%) 1331 BC 1326 (23.8%) 1291 BC 1279 (4.8%) 1271 BC	1409 (92.7%) 1258 BC 1233 (2.7%) 1218 BC
130416/80	Poz-57382	No tomb attribution, Wupu	Wood/poplar trunk from the tomb chamber	2715 ± 35	896 (68.2%) 829 BC	924 (95.4%) 806 BC
130417/82	Poz-76927	Yanghai, M195	Leather fragment/bow and arrow sheath	2475 ± 30	753 (19.4%) 702 BC 696 (5.6%) 682 BC 669 (22.3%) 611 BC 594 (20.9%) 540 BC	771 (93.6%) 477 BC 463 (0.6%) 456 BC 445 (1.3%) 431 BC
130417/86	Poz-76845	Yanghai, M195	Leather fragment/bow and arrow sheath	3400 ± 170 very poor date, not used	1916 (68.2%) 1503 BC	2197 (0.8%) 2168 BC 2149 (92.9%) 1371 BC 1359 (1.8%) 1299 BC

construct median-joining haplotype networks (Bandelt et al., 1999) using PopART 1.7 (Leigh, n.d.).

Results

Ancient samples

Sequences were recovered in five samples from Wupu and four samples from Yanghai (Table 1). One sample from Yanghai did not yield PCR products. In all successful amplifications from Wupu, both species and control region haplotype were recovered, while in the successful Yanghai amplifications the mitochondrial haplotype could only be determined in a single sample.

The three samples of the Wupu leather coat 86HWM:1-00751 were identified as sheep skin. One (130414/24) belongs to sheep haplogroup A while 130414/28 belongs to haplogroup

B; 130414/26 shows intermediate characteristics between haplogroups A and B. Only one sample from Yanghai, the bracer 2003SAYIM195:12 (130417/87), was amplifiable as a sheep control regional fragment (fragment 1), which corresponds to haplotype A1a.

Both samples from the Wupu leather coat, 86HWM:2-00752, were identified as goat-derived and belonged to goat haplogroup A, with 130414/33 being closely related to haplotype A2 and 130414/34 to A3.

The first cytB fragment of 130417/84 shows two transitions that are not known from modern *Bos taurus* haplotypes and may not represent a domestic animal, but a wild aurochs or yak (*Bos grunniens*). Unfortunately the control region sequence has not been amplified successfully so far because of poor DNA quality.

Table 3. PCR primers used in this study.

Primer ID	Sequence 5'–3'	Length	Source
cytB			
142f	GGCCTATTCCTAGCRATACAC	21	This study
248r	TGTATRTATCGGATRATTTCAGC	22	This study
395f	TCYTACCATGAGGACAAATATC	22	This study
500r	CCDCCTCAGATYCATTTCGAC	20	This study
931f	AARCAACGVAGCATRATATCCCG	23	This study
1022r	GMCCTCCRATTCATGTRAGTG	21	This study
Ovis ctr			
LI5391	CCACTATCAACACCCAAAG	19	Cai et al. (2011)
HI5534	AAGTCCGTGTTGTATGTTTG	20	Cai et al. (2011)
OAU15993	GCATGTAGGGTATTAACCTGCTTGAC	26	Geörg (2013)
OAL16087	GATCCTTGCRYAGCGGGTTG	20	Geörg (2013)
OAU16068	CCAYTAGATCACGAGCTTGTTCCAC	24	Geörg (2013)
OAL16161	CTGAAGAAAGAACCAGATGCCTGT	24	Geörg (2013)
Capra ctr			
CAP-FII	GATCTTCCYCATGCATATAAGCA	23	Fernández et al. (2006)
CAP-RII	CGGGTTGCTGGTTTCAC	17	Fernández et al. (2006)
Bos ctr			
An1F	CTTAATTACCATGCCGCGTG	20	Cai et al. (2014)
An3R	CGAGATGTCTTATTTAAGAGG	21	Cai et al. (2014)

Modern reference samples

Control region sequences were recovered from seven sheep and three goats from Ya'erguolecun, China. Three sheep from one farm possessed haplotype A1c, one sheep from a second farm belonged to haplogroup A, while the other three belonged to haplogroup B. Two goats belonged to haplogroup A3, while one goat belonged to haplogroup A2.

Discussion

All successfully identified leather samples were derived from domestic animals. Neither wild sheep (argali *Ovis ammon*) nor wild goats (Siberian ibex *Capra sibirica*), which are both native to Central Asia, were found. The results show that all common domesticated bovines were used to produce leather in the sites of Wupu and Yanghai. Sheep, goat, and cattle were present in central and eastern Asia at least 5000 years ago (Cai et al., 2011, 2014; Lv et al., 2015; Payne and Hodges, 1997; Zhang et al., 2013).

Unfortunately, even after several repeats, mitochondrial haplotypes were not determined for most samples from the Yanghai site. This may be a result of different taphonomic conditions, which are less conducive for ancient DNA preservation, compared with Wupu or a result of different storage conditions in the respective museum collections, which have a significant influence on the success rate of ancient DNA analyses (Pruvost et al., 2007). Furthermore, among the leather objects from Wupu (coats), different tanning methods may have been employed compared with the objects from Yanghai (quivers and bracer).

Sheep

According to data from Lv et al. (2015), the first sheep entered eastern and northern China from the Caucasus, via Central Asia and the Mongolian Plateau around 6800–6400 years ago, with a second migratory wave around 4500 years ago. Today, the most common sheep mitochondrial haplogroups are A and B, with A dominating in eastern and southern Asia, while B is the most common haplogroup in Europe and western Asia (Lv et al., 2015; Pedrosa et al., 2005; Tapio et al., 2006). Haplogroup C is common in fat-tailed breeds and may have entered China during the second migratory wave (Lv et al., 2015) while D and E are only found in

southwest Asia (Tapio et al., 2006). Previously, this pattern was also found in an ancient DNA study (Cai et al., 2011) of east Chinese sheep remains from c. 4500 to 1500 years ago, where A was the predominant haplogroup at 95.5%.

The three sheep samples from Wupu belonged to haplogroups A, B, and an intermediate haplotype between the two, respectively. The control region sequence of /24 (A) has only one mutation difference with the most common A haplotype (Figure 3) which is common in modern Chinese sheep and was also found in one of the modern Turpan sheep. The control region sequence of /28 (B) was, overall, the most common in our dataset and is found in Europe, the Near East, and China (Figure 3). Three modern Turpan sheep shared the same sequence. The unusual intermediate control region sequence of /26 was previously found in modern sheep from Pakistan (JX235856) and Italy (JN184935) (Pariset et al., 2011) but is unknown from modern China.

Our findings show that a certain degree of diversity was present in domesticated sheep from the Wupu site, as both haplogroups A and B were present in a small sample. Haplogroup C was not detected, which could imply that the sheep from which the leather was derived did not belong to the ancestors of the modern fat-tailed breeds (Lv et al., 2015). However, none of the sampled modern Turpan fat-tailed sheep showed this haplotype either, so this prediction is questionable. The similarities of the control region sequences between ancient samples and modern Turpan and other Chinese breeds indicate a continuity of sheep husbandry in the Turpan Oasis from the 1st millennium BC to modern times.

Goat

Most likely, the first domesticated goats entered eastern Asia at the same time as sheep. Modern goat mitochondrial haplotype diversity is dominated by haplogroup A with a global preponderance of over 90% (Luikart et al., 2001; Naderi et al., 2007), which may correspond to the first wave of migration of early domestics from the center of origin of domestication in the Near East (Akis et al., 2014; Luikart et al., 2001). In eastern Asia, haplogroup B is also present in moderate numbers (~25%) (Chen et al., 2005; Liu et al., 2009) and was found in ancient European and Chinese goats (Fernández et al., 2006; Han et al., 2010) while haplogroups C–D are rare worldwide. In this context, it is not surprising that




Figure 3. Median-joining haplotype network of the control region fragments of 58 ancient and modern sheep. Ancient Chinese sheep denote results from leather samples of this study. Turpan sheep are modern sheep sampled from two farms close to the archaeological sites of this study.




Figure 4. Median-joining haplotype network of the amplified control region fragments of 62 ancient and modern goats. Ancient Chinese goats denote results from leather samples of this study. Turpan goats are modern goats sampled from two farms close to the archaeological site of this study.

both goat-derived leather samples from Wupu belonged to the most common haplogroup A. Both sequences were previously found in modern Chinese domesticated goats (/33:DQ089113, DQ089147, and /34:DQ089186). Accordingly, modern goats from Turpan also belonged to this haplogroup with two individuals being particularly close to /34, each differing by only one transition (Figure 4). As in sheep this implies a continuity of goat farming in the Turpan Oasis.

Conclusion

Our results show that domestic sheep, goat, and cattle were present and used for leather production in the Turpan and Hami oases around *c.* 2800–2500 years ago. Based on the directly dated cheek-piece from Wupu, the domesticated horse and horse riding in the Hami region can be dated to *c.* 1100–900 BC. The mitochondrial control region sequences of ancient sheep and goats are identical or closely related to modern domesticated animals from the same geographic region and other parts of China, pointing to a continuity of pastoralism with the same general stock of animals to the present. Wild ungulates apparently did not play an important role for the production of leather garments and gear in the Bronze Age communities of Wupu and Yanghai, emphasizing the importance of animal husbandry. Our results demonstrate that ancient DNA methods are valuable for identifying species and lineages of Central Asian domesticated animals from archaeological leather remains.

Acknowledgements

The authors acknowledge gratefully the work of Chinese colleagues who excavated and preserved the objects in Turpan and Hami and who shared in the discussion of the archaeological context. Thanks to P Wertmann, XC Chen, and D Hosner for their help in organizing fieldwork campaigns, providing photos, and translation from Chinese and to D Lieckfeldt for assisting in lab work and primer design. We also thank Dr R Spengler for editorial corrections and valuable suggestions and two anonymous reviewers for their helpful comments which we considered in the revised version of this manuscript.

Funding

This study is a contribution to the ‘Silk Road Fashion’ research project supported by the German Federal Ministry of Research and Education (Grant 01UO1310) and the German Archaeological Institute.

References

- Akis I, Onar V, Toker N et al. (2014) Ancient DNA analysis of Anatolian goat remains excavated from a Urartian castle in eastern Turkey. *International Journal of Osteoarchaeology*. Epub ahead of print 25 August. DOI: 10.1002/oa.2415.
- Bandelt HJ, Forster P and Röhl A (1999) Median-joining networks for inferring intraspecific phylogenies. *Molecular Biology and Evolution* 16(1): 37–48.
- Beck U, Wagner M, Li X et al. (2014) The invention of trousers and its likely affiliation with horseback riding and mobility: A case study of late 2nd millennium BC finds from Turfan in eastern Central Asia (The Bridging Eurasia Research Initiative: Modes of mobility and sustainability in the palaeoenvironmental and archaeological archives from Eurasia). *Quaternary International* 348: 224–235.
- Brandt LØ, Tranekjer LD, Mannering U et al. (2011) Characterising the potential of sheep wool for ancient DNA analyses. *Archaeological and Anthropological Sciences* 3(2): 209–221.
- Bronk Ramsey C (2013) *OxCal Program v4.2.4*. Oxford: Radiocarbon Accelerator Unit, University of Oxford.
- Cai D, Sun Y, Tang Z et al. (2014) The origins of Chinese domestic cattle as revealed by ancient DNA analysis. *Journal of Archaeological Science* 41: 423–434.
- Cai D, Tang Z, Yu H et al. (2011) Early history of Chinese domestic sheep indicated by ancient DNA analysis of Bronze Age individuals. *Journal of Archaeological Science* 38(4): 896–902.
- Chen SY, Su YH, Wu SF et al. (2005) Mitochondrial diversity and phylogeographic structure of Chinese domestic goats. *Molecular Biology and Evolution* 37: 804–814.
- Cooper A and Poinar HN (2000) Ancient DNA: Do it right or not at all. *Science* 289(5482): 1139.
- Dabney J, Knapp M, Glocke I et al. (2013) Complete mitochondrial genome sequence of a Middle Pleistocene cave bear reconstructed from ultrashort DNA fragments. *Proceedings of the National Academy of Sciences of the United States of America* 110(39): 15758–15763.
- Domrös M and Peng G (1988) *The climate of China*. Berlin: Springer.
- Dong G-H, Wang Z-L, Ren L-L et al. (2014) A comparative study of ¹⁴C dating on charcoal and charred seeds from Late Neolithic and Bronze Age sites in Gansu and Qinghai provinces, NW China. *Radiocarbon* 56(1): 157–163.
- Fernández H, Hughes S, Vigne J-D et al. (2006) Divergent mtDNA lineages of goats in an Early Neolithic site, far from the initial domestication areas. *Proceedings of the National Academy of Sciences of the United States of America* 103(42): 15375–15379.
- Geörg C (2013) *Paläopopulationsgenetik von Schwein und Schaf in Südosteuropa und Transkaukasien*. Dissertation, German Archaeological Institute.
- Guojia Wenwuju (2012) *Zhongguo wen wu di tu ji: Xinjiang Weiwu'er zizhiqu fen ce* [Atlas of Chinese Cultural Relics: Xinjiang Uyghur Autonomous Region Volume]. Beijing: Cultural Relics Publishing House.
- Han L, Yu H-X, Cai D-W et al. (2010) Mitochondrial DNA analysis provides new insights into the origin of the Chinese domestic goat. *Small Ruminant Research* 90(1–3): 41–46.
- Hofreiter M, Paijmans JLA, Goodchild H et al. (2015) The future of ancient DNA: Technical advances and conceptual shifts. *Bioessays* 37: 284–293.
- Jiang H-E, Li X, Ferguson DK et al. (2007) The discovery of *Capparis spinosa* L. (Capparidaceae) in the Yanghai Tombs (2800 years B.P.), NW China, and its medicinal implications. *Journal of Ethnopharmacology* 113: 409–420.
- Jiang H-E, Li X, Zhao Y-X et al. (2006) A new insight into *Cannabis sativa* (Cannabaceae) utilization from 2500-year-old Yanghai Tombs, Xinjiang, China. *Journal of Ethnopharmacology* 108: 414–422.
- Jiang H-E, Zhang Y-B, Li X et al. (2009) Evidence for early viticulture in China: Proof of a grapevine (*Vitis vinifera* L., Vitaceae) in the Yanghai Tombs, Xinjiang. *Journal of Archaeological Science* 36(7): 1458–1465.
- Kramell A, Li X, Csuk R et al. (2014) Dyes of late Bronze Age textile clothes and accessories from the Yanghai archaeological site, Turfan, China: Determination of the fibers, color analysis and dating (The Bridging Eurasia Research Initiative: Modes of mobility and sustainability in the palaeoenvironmental and archaeological archives from Eurasia). *Quaternary International* 348: 214–223.
- Leigh J (n.d.) PopART website. Available at: <http://popart.otago.ac.nz> (accessed 8 December 2015).
- Li C, Li H, Cui Y et al. (2010) Evidence that a West-East admixed population lived in the Tarim Basin as early as the early Bronze Age. *BMC Biology* 8: 15.
- Li C, Lister DL, Li H et al. (2011) Ancient DNA analysis of desiccated wheat grains excavated from a Bronze Age cemetery in Xinjiang. *Journal of Archaeological Science* 38: 115–119.

- Liu Y-P, Cao S-X, Chen S-Y et al. (2009) Genetic diversity of Chinese domestic goat based on the mitochondrial DNA sequence variation. *Journal of Animal Breeding and Genetics* 126: 80–89.
- Luikart G, Gielly L, Excoffier L et al. (2001) Multiple maternal origins and weak phylogeographic structure in domestic goats. *Proceedings of the National Academy of Sciences of the United States of America* 98: 5927–5932.
- Lv F-H, Peng W-F, Yang J et al. (2015) Mitogenomic meta-analysis identifies two phases of migration in the history of eastern Eurasian sheep. *Molecular Biology and Evolution* 32: 2515–2533.
- Naderi S, Rezaei H-R, Taberlet P et al. (2007) Large-scale mitochondrial DNA analysis of the domestic goat reveals six haplogroups with high diversity. *PLoS ONE* 2(10): e1012.
- Pariset L, Mariotti M, Gargani M et al. (2011) Genetic diversity of sheep breeds from Albania, Greece, and Italy assessed by mitochondrial DNA and nuclear polymorphisms (SNPs). *The Scientific World Journal* 11: 1641–1659.
- Payne WJA and Hodges J (1997) *Tropical Cattle: Origins, Breeds and Breeding Policies*. Oxford; Malden, MA: Blackwell Science Ltd.
- Pedrosa S, Uzun M, Arranz J-J et al. (2005) Evidence of three maternal lineages in near eastern sheep supporting multiple domestication events. *Proceedings of the Royal Society of London B: Biological Sciences* 272(1577): 2211–2217.
- Peng M-S, Fan L, Shi N-N et al. (2015) DomeTree: A canonical toolkit for mitochondrial DNA analyses in domesticated animals. *Molecular Ecology Resources* 15(5): 1238–1242.
- Poulakakis N, Tselikas A, Bitsakis I et al. (2007) Ancient DNA and the genetic signature of ancient Greek manuscripts. *Journal of Archaeological Science* 34(5): 675–680.
- Pruvost M, Schwarz R, Correia VB et al. (2007) Freshly excavated fossil bones are best for amplification of ancient DNA. *Proceedings of the National Academy of Sciences of the United States of America* 104(3): 739–744.
- Reimer PJ, Bard E, Bayliss A et al. (2013) IntCal13 and Marine13 radiocarbon age calibration curves 0–50,000 years cal BP. *Radiocarbon* 55(4): 1869–1887.
- Schlumbaum A, Campos PF, Volken S et al. (2010) Ancient DNA, a Neolithic legging from the Swiss Alps and the early history of goat. *Journal of Archaeological Science* 37(6): 1247–1251.
- Tapio M, Marzanov N, Ozerov M et al. (2006) Sheep mitochondrial DNA variation in European, Caucasian, and Central Asian areas. *Molecular Biology and Evolution* 23(9): 1776–1783.
- Teasdale MD, van Doorn NL, Fiddymont S et al. (2015) Paging through history: Parchment as a reservoir of ancient DNA for next generation sequencing. *Philosophical Transactions of the Royal Society of London Series B: Biological Sciences* 370(1660): 20130379.
- Vuissoz A, Worobey M, Odegaard N et al. (2007) The survival of PCR-amplifiable DNA in cow leather. *Journal of Archaeological Science* 34(5): 823–829.
- Wagner M, Bo W, Tarasov P et al. (2009) The life-history of the warrior-and-centaur trousers from Sampula (Xinjiang, China). *Antiquity* 83: 1065–1075.
- Wagner M, Wu X, Tarasov P et al. (2011) Radiocarbon-dated archaeological record of early first millennium B.C. mounted pastoralists in the Kunlun Mountains, China. *Proceedings of the National Academy of Sciences of the United States of America* 108(38): 15733–15738.
- Wang BH (1999) *Xinjiang gu shi* [The ancient corpses of Xinjiang]. Ürümqi: Xinjiang Renmin Chubanshe.
- Xinjiang tu lu fan xue yan jiu yuan and Xinjiang wen wu kao gu yan jiu suo (2011) Xinjiang Shanshan Yanghai mu di fa jue bao gao. *Kaogu Xuebao* 1: 99–150.
- Zeder MA (2008) Domestication and early agriculture in the Mediterranean Basin: Origins, diffusion, and impact. *Proceedings of the National Academy of Sciences of the United States of America*. Epub ahead of print 27 May. DOI: 10.1073/pnas.0801317105.
- Zhang H, Paijmans JLA, Chang F et al. (2013) Morphological and genetic evidence for early Holocene cattle management in northeastern China. *Nature Communications* 4: 2755.