

# Quantitatively evaluating the effects of climate factors on runoff change for Aksu River in northwestern China

Baofu Li · Yaning Chen · Heigang Xiong

Received: 5 July 2014 / Accepted: 8 December 2014  
© Springer-Verlag Wien 2014

**Abstract** Much attention has recently been focused on the effects that precipitation and potential evapotranspiration (PET) have had on runoff change; however, the influence of temperature on runoff needs to be further studied. We attempted to employ the improved elasticity method to evaluate the effects of climate factors (CF, especially temperature) on runoff change for Aksu River in the arid region of northwest China. Data from Aksu River in the arid region of northwest China were analyzed to investigate changes in annual runoff and CF during the period of 1960–2010. The key findings of this study indicated that the annual runoff had a significant ( $P < 0.01$ ) increasing trend with a rate of  $3.78 \times 10^8 \text{ m}^3/\text{decade}$ , and the temperature and precipitation also exhibited significant rising trends, at a rate of  $0.28 \text{ }^\circ\text{C}/\text{decade}$  ( $P < 0.01$ ) and  $15.11 \text{ mm}/\text{decade}$  ( $P < 0.05$ ), respectively, while PET showed a decreasing trend ( $22.66 \text{ mm}/\text{decade}$ ,  $P < 0.01$ ). Step change point in runoff occurred in the year 1993. Thus, we employed the mean runoff and climate factors during the period 1960–1993 as the benchmark value to measure the change. In 1994–2010, mean runoff increased by 22 %. Results also revealed that temperature rising was the most important factor that increased runoff with contribution of 45 %, while precipitation and PET were responsible for 22 and 27 % of the runoff change, respectively, indicating that the runoff of

increasing percentage only accounted for 6 % owing to human activities and other factors, and showed that climate variability was the main reason for the runoff change in Aksu River.

## 1 Introduction

Climate variation has made the global water cycle remarkably change (Milly et al. 2005; Chen et al. 2013; Chen and Chen 2014). With a worsening of the water shortage problem and increasing of global average temperature, much attention has recently been focused on the effects of climate variation on hydrological cycle (IPCC 2007). As a particular case, the air temperature in the arid region of northwest China has been increasing by a rate of  $0.33\text{--}0.39 \text{ }^\circ\text{C}/\text{decade}$  in recent 50 years (Zhang et al. 2010; Li et al. 2012a, 2013a), higher than the average of China ( $0.25 \text{ }^\circ\text{C}/\text{decade}$ ) (Ren et al. 2005) and that of the entire globe ( $0.13 \text{ }^\circ\text{C}/\text{decade}$ ) (IPCC 2007) for the same period. This further exacerbated the vulnerability and uncertainty of water resources system that is mainly recharged by mountain glacier and snow melt water (Xu et al. 2011). Therefore, quantitatively evaluating these effects is important for regional water resources assessment and management.

Recently, some new approaches have been proposed to explore the effects of climate change and human activities on runoff (Ma et al. 2008a; Liu et al. 2010; Zhang et al. 2012; Meng and Mo 2012; Jiang et al. 2012; Li et al. 2014a). For example, Jiang et al. (2011) have identified the climate variability and human activities on runoff from the Laohahe Basin by using double cumulative curve method. Zheng et al. (2009) attempted to analyze the responses of streamflow to precipitation, potential evapotranspiration (PET), and land surface change in the headwaters of the Yellow River Basin. Meanwhile, hydrologic sensitivity analysis method has been employed to probe into the effects of precipitation and PET on runoff for Kaidu River Basin in arid region of northwest China

---

B. Li (✉)  
College of Geography and Tourism, Qufu Normal University,  
Rizhao 276826, China  
e-mail: lenny006@163.com

B. Li · Y. Chen  
State Key Laboratory of Desert and Oasis Ecology, Xinjiang Institute  
of Ecology and Geography, Chinese Academy of Sciences,  
Urumqi 830011, China

H. Xiong  
College of Applied Arts and Science, Beijing Union University,  
Beijing 100083, China

(Chen et al. 2013) and to quantitatively assess the impacts of climate variability and human activities on runoff changes in Haihe River Basin (Wang et al. 2013). Zang et al. (2013) illustrated the impacts of human activities and climate variability on green and blue water flows in the Heihe River Basin in northwest China. Ye et al. (2009, 2013) distinguished the relative impacts of climate change and human activities on variation of streamflow in the Poyang Lake catchment in China by using a coupled water and energy budgets analysis. Li et al. (2014b) utilized statistical methods to separate the impacts of climate variation and human activities on runoff in the Songhua River Basin of northeast China. Zeng et al. (2014) investigated the impacts of climate on runoff changes at annual, seasonal, and monthly time scales in the Zhang River Basin of north China-based hydrological modeling and sensitivity method. It can be seen that more research focus in humid and semi-humid regions, but less study has been done on special hydrological processes in arid areas. Moreover, several selected methods, such as gray correlation analysis method (Yang et al. 2005), linear regression method (Jiang et al. 2011), the double mass curve method (Jiang et al. 2012), water balance method, hydrological sensitivity analysis (Chen et al. 2013; Ye et al. 2009, 2013; Li et al. 2014b; Zeng et al. 2014), hydrological model (Zhang et al. 2012; Zang et al. 2013; Li et al. 2014a), slope change ratio of accumulative quantity method (Wang et al. 2012), etc., were applied to quantitatively differentiate the effects of climate change and human activities on runoff. Hydrological model method has a strong physical basis, but there is still obvious uncertainty (Song et al. 2013). Statistical analysis method can be easily used and demand detailed long-term period observation hydrologic and meteorological data for the basin. These methods can be adopted to quantitatively indicate the impacts of precipitation and PET on runoff (Ye et al. 2009, 2013; Li et al. 2014b); moreover, the effect of temperature on runoff can be only reflected by PET (Zheng et al. 2009; Chen et al. 2013; Wang et al. 2013; Ye et al. 2013; Chen and Chen 2014). However, with the intensifying global warming, the influence of temperature change on water resources gradually strengthened, especially the surface runoff recharges mainly from glacier and snow melt water in arid region rivers, which cannot be reflected accurately the impact of climate change on runoff only by PET. Therefore, these methods are so limited that it is difficult to calculate runoff change directly generated by temperature rising (Chen et al. 2013; Chen and Chen 2014). Thus, it is important to understand the hydrological process responses to different climate factor (temperature, precipitation, and PET) changes in order to develop sustainable basin management strategies.

Aksu River is the main tributary of the Tarim River, and its upper reaches are in the Kunlun Mountains, nearly half of the water from the mountains of glacier and snow melt water in the Tarim Basin. Some scholars have conducted research on

the correlation between Aksu River runoff and regional climate change in recent years. Jiang et al. (2005) analyzed both intra-annual distribution law and variation characteristics over years of different supply sources runoff and come to conclusion that annual runoff discharge from glacier and snow smelt water has been increasing since 1990 in mountain area and the rise of temperature has a huge effect on runoff increase than that of precipitation in Aksu River. Li et al. (2008) investigated the relationship between North Atlantic Oscillation and Aksu River runoff by using the methods of wavelet transform and cross wavelet transform and indicated the atmospheric circulation variations caused by NAO has an impact on the climate of the Aksu River Basin; meanwhile, the runoff of the Aksu River is affected. Chen et al. (2009) pointed out that the increase of temperature in the tributaries of the Aksu River is higher than that in the tributaries of the Yarkand River and Hotan River in the Tarim Basin. Yu et al. (2011) checked the nonlinear characteristics of annual runoff processes from 1957 to 2008 and illustrated that the annual runoff in Aksu River will show an increasing trend in the future. Xu et al. (2011) found that there was a close relationship between variations in the annual runoff of the Aksu River and regional climate change. Li et al. (2012b) found that if runoff recharge proportion from glacier and snow smelt water is very large, the correlation between runoff and temperature is significantly positive in typical river area. Predecessors' research results show that annual runoff discharge from glacier and snow smelt water is about 42 % (Jiang et al. 2005); thus, we guess that the influence of temperature on runoff is very important in the Aksu River. These studies analyzed qualitatively the influence of climate on Aksu River runoff but did not explore quantitatively the effect of climate factors on runoff.


In this study, firstly, we make an attempt to improve elasticity method based on the principle of elasticity theory; secondly, we further quantitatively evaluate the impact of temperature, precipitation, and PET on annual runoff change for Aksu River in the arid region of northwest China by using the improved elasticity method.

## 2 Materials and methods

### 2.1 Study area

Located on the south slope of the Tianshan Mountains and the northwest edge of the Tarim Basin, Aksu River is the largest runoff of the rivers and is enclosed between latitudes 40° 17'–42° 27' N and longitudes 75° 35'–80° 59' E (Fig. 1) with a basin area of  $5.14 \times 10^4$  km<sup>2</sup> (Xu et al. 2011). The topography descends gradually from north to south and from west to east, featured by unique terrain from high to low, low-middle mountain and hill, group of piedmont pluvial fans, tilted

**Fig. 1** Location of the Aksu River and the distribution of meteorological station and hydrological station



alluvial-proluvial plain, and alluvial plain (elevation of 1000–1500 m) in turn. The Aksu River belongs to temperate continental arid climate, which is characterized by drought, lack of rainfall, intensity evaporation, and large temperature difference between days and years, because of the distinct geographical position and far away from sea; the drainage's average annual temperature, precipitation, and PET are 9.2–11.5 °C, 64 mm, and 1890 mm, respectively (Xu et al. 2011); annual extreme maximum and minimum temperatures are 40.2 and −27.6 °C, respectively, and the mean annual sunshine duration is 2850 h. The Kumalak and Toxkan River, as two main tributaries, join at Kaladuwei before flowing into the Aksu River. Mountainous areas are the major areas of runoff generation for the Aksu River, and the complex climatic condition and hydrological environment causes the runoff fluctuation.

## 2.2 Data

Monthly runoff data from the Xiehela and Shaliguilake hydrological station in the mountains-pass, which was available for the period from 1960 to 2010, were used in this study. To reflect the effect of mountain climate on runoff, we selected the meteorological station (Aheqi and Toergate) to represent climate variations of mountains area (average elevation of 2746 m). Daily relative humidity, maximum and minimum air temperature, wind speed, and sunshine hours from meteorological station during the period of 1960–2010 were utilized to calculate potential evapotranspiration by the Penman-Monteith equation recommended by FAO (Allen et al. 1998).

All meteorological station data selected for this study had been maintained following the standard methods of the National Meteorological Administration of China, having strict high-quality data control (including extreme inspection and time consistency checks) before releasing these data. The

runoff data in each river were obtained from local Hydrology Bureaus. Meanwhile, using the RCLimDex software package (available at the ETCCDI website, <http://cccma.seos.uvic.ca/ETCCDI/software.shtml>) attained data quality control and homogeneity assessment (You et al. 2011).

## 2.3 Method

### 2.3.1 Elasticity theory method

The concept of elasticity theory could be used to detect the effect of an independent variable on the dependent variable (Jin and Wu 2002). To quantitatively evaluate the impacts of climate factors on the runoff change, we first assume that the attributions of runoff change to climate factors (CF) and non-CF-related change (such as human activities and other factors) can be approximated as follows:

$$\Delta Q = f'_C \Delta C + \Delta f'_N \Delta N \quad (1)$$

$$f'_C \Delta C = f'_T \Delta T + f'_P \Delta P + f'_E \Delta E \quad (2)$$

where  $\Delta Q$ ,  $\Delta C$ , and  $\Delta N$  are changes in runoff, CF (including  $\Delta T$ ,  $\Delta P$ , and  $\Delta E$  are changes in temperature, precipitation, and potential evapotranspiration, respectively) and non-CF, respectively, with  $f'_T = \partial Q / \partial T$ ,  $f'_P = \partial Q / \partial P$ ,  $f'_E = \partial Q / \partial E$ , and  $f'_N = \partial Q / \partial N$ . If we assume that the every factor is independent of other change factors, Eqs. (1) and (2) can be written as the following:

$$\Delta Q = \Delta Q_T + \Delta Q_P + \Delta Q_E + \Delta Q_N \quad (3)$$

$$\Delta Q_T = f'_T \Delta T \quad (4)$$

$$\Delta Q_P = f'_P \Delta P \tag{5}$$

$$\Delta Q_E = f'_E \Delta E \tag{6}$$

$$\Delta Q_N = f'_N \Delta N \tag{7}$$

where  $\Delta Q_T$ ,  $\Delta Q_P$ ,  $\Delta Q_E$ , and  $\Delta Q_N$  change in runoff due to temperature, precipitation, potential evapotranspiration, and non-CF, respectively. This framework is used to separate the effects of CF and non-CF factors on runoff.

Following the method (Schaake 1990), this study defines the climate factor ( $C_i$ ) elasticity regarding runoff ( $\varepsilon_i$ ) as the ratio of the proportional change of runoff to the proportional change of the factor:

$$\varepsilon_i = \frac{dQ/Q}{dC_i/C_i} \tag{8}$$

$i \in \{\text{temperature, precipitation, potential evapotranspiration}\}$

The physical meaning of  $\varepsilon_i$  is that 1 % of a factor change may lead to  $\varepsilon_i\%$  of runoff change. Based on Eq. (8), we rewrite Eq. (4–6) as

$$\Delta Q_i = (\varepsilon_i \Delta C_i / C_i) \cdot Q \tag{9}$$

$i \in \{\text{temperature, precipitation, potential evapotranspiration}\}$

From Eq. (8) we also see

$$\varepsilon_i = \frac{dQ}{dC_i} \frac{C_i}{Q} = f'_{C_i} C_i / Q \tag{10}$$

$i \in \{\text{temperature, precipitation, potential evapotranspiration}\}$

If the relationship between runoff and CF is known, the CF elasticity can be derived mathematically.

Following the literature (Schaake 1990; Sankarasubramanian et al. 2001; Zheng et al. 2009), we utilize a nonparametric estimator to describe the relationship between runoff and CF directly based on observation values:

$$(Q_j - Q_b) / Q_b = \varepsilon_i (C_{ij} - C_{ib}) / C_{ib} \tag{11}$$

where  $\varepsilon_i$  is elasticity of runoff with respect to CF;  $Q_j$  and  $C_{ij}$  are yearly runoff and the factor values, with  $j=1960, 1961, \dots, 2010$ ;  $Q_b$  and  $C_{ib}$  are the benchmark value for the runoff and CF, respectively. Therefore,  $\varepsilon_i$  can be regarded as the linear regression coefficient between  $(Q_j - Q_b) / Q_b$  and  $(C_{ij} - C_{ib}) / C_{ib}$ .

To quantitatively identify runoff change caused by CF change in the past 50 years, the following formula is used to

calculate the change rate of runoff:

$$\varphi_i = 100\% \times \left( \varepsilon_i \cdot \frac{\Delta C_i}{C_{ib}} \cdot Q_b \right) / \Delta Q \tag{12}$$

where  $\Delta C_i$  and  $\Delta Q$  are the changes of CF and runoff during the period 1960–2010, respectively. Thus,  $\varphi_i$  is the importance of the CF variation in runoff change in recent 50 years.

A vast number of climate elastic analysis methods exist; however, the elastic analysis based on nonparametric method has higher effectiveness and stability (Sankarasubramanian et al. 2001). To verify the result of this paper, we adopt the nonparametric elastic analysis method proposed by Zheng et al. (2009) to explore the sensitivity of annual runoff to the CF. The method is widely utilized to present the sensitivity of hydrological elements to changes in meteorological elements in the process of the effect of climate change on hydrological system (Zheng et al. 2009; Chen et al. 2012; Li et al. 2013b). The formula is as follows (Zheng et al. 2009):

$$\varepsilon = \frac{\bar{X} \sum (X_i - \bar{X})(Q_i - \bar{Q})}{\bar{Q} \sum (X_i - \bar{X})^2} \tag{13}$$

where  $X_i$  denotes the meteorological element,  $Q_i$  denotes the annual runoff,  $\varepsilon$  is the sensitivity coefficient, and  $\bar{X}$  and  $\bar{Q}$  are mean values of runoff and meteorological element in many years, respectively.

### 2.3.2 Trend test

We employed the Mann-Kendall (MK) statistical test (Mann 1945; Kendall 1975) to test the significance of trends in the annual runoff, temperature, and PET in the study area. The nonparametric Mann-Kendall statistical test has been commonly adopted to assess the significance of monotonic trends in meteorological and hydrologic series (Chen and Xu 2005; Zhang et al. 2011). For a time series  $X = \{x_1, x_2, \dots, x_n\}$ , when  $n > 10$ , the standard normal statistic  $Z$  is estimated as follows:

$$Z = \begin{cases} (S-1)/\sqrt{\text{var}(S)} & S > 0 \\ 0 & 0 \\ (S+1)/\sqrt{\text{var}(S)} & S < 0 \end{cases} \tag{14}$$

where

$$S = \sum_{i=1}^{n-1} \sum_{j=i+1}^n \text{sgn}(x_j - x_i) \tag{15}$$

$$\text{sgn}(\theta) = \begin{cases} +1, & \theta > 0 \\ 0, & \theta = 0 \\ -1, & \theta < 0 \end{cases} \quad (16)$$

$$\text{var}(S) = \left[ n(n-1)(2n+5) - \sum_t t(t-1)(2t+5) \right] / 18 \quad (17)$$

where  $t$  is the extent of any given tie, and  $\sum_t$  denotes the summation of all ties.

### 2.3.3 Step change point analysis

The nonparametric Mann-Kendall-Sneyers test (Mann 1945; Kendall 1975; Sneyers 1975) was applied in this study for determining the occurrence of step change points of climate factor and runoff.  $x_1, \dots, x_n$  represent the data points. The numbers  $m_i$  of elements  $x_j$  preceding it ( $j < i$ ) such that  $x_j < x_i$  are computed for each element  $x_i$ . Under the null hypothesis (no step change point), the normally distributed statistic  $t_k$  can be described as follows:

$$t_k = \sum_{i=1}^k m_i \quad (2 \leq k \leq n) \quad (18)$$

$t_k$  as mean and variance of the normally distributed statistic can be calculated as follows:

$$\bar{t}_k = E(t_k) = k(k-1)/4 \quad (19)$$

$$\text{var}(t_k) = k(k-1)(2k+5)/72 \quad (20)$$

$u_k$  as the normalized variable statistic is given in following formula:

$$u_k = (t_k - \bar{t}_k) / \sqrt{\text{var}(t_k)} \quad (21)$$

## 3 Results

### 3.1 Trend of CF and runoff

Figure 2 shows long-term variations in annual runoff, temperature, precipitation, and potential evapotranspiration (PET) for Aksu River from 1960 to 2010. Mann-Kendall statistical test reveals that the runoff and temperature exhibits an increasing

trend at  $P < 0.01$  significant level, with the rate of  $3.78 \times 10^8$   $\text{m}^3/\text{decade}$  and  $0.28$   $^\circ\text{C}/\text{decade}$ , respectively. Precipitation shows significant upward trend ( $P < 0.05$ ) with the rate of  $15.11$   $\text{mm}/\text{decade}$ . However, the PET has a significant decreasing trend ( $P < 0.01$ ) at a rate of  $22.66$   $\text{mm}/\text{decade}$ .

The Mann-Kendall-Sneyers test was applied to detect the step change point of the annual runoff during 1960 to 2010. Figure 3 shows the computed probability series of the step change point years for runoff. The intersection of the curves indicates that there is a step change point in 1993 (at  $P < 0.05$  significance level) for the runoff. The value is consistent with the results from previous research (Li et al. 2008). To investigate the effect of CF on runoff, we also carried out the test for annual temperature, precipitation, and PET. The results reveal that abrupt changes in temperature (1993), precipitation (1992 and 1994), and PET (1990) basically occurred in the early 1990s. That is to say, from 1994 to 2010, runoff had significant change caused by climate and other factors. In this study, we used the mean runoff and CF of 1960–1993 as the benchmark value to measure the change. Concurrently, annual runoff, temperature, and precipitation in 1994–2010 was more about 22, 60, and 23 %, respectively, than that of the period 1960–1993, while annual PET was less 5 %.


### 3.2 CF elasticity of runoff

Figure 4 shows the linear regression relationships between the proportional change in temperature, precipitation, PET, and that in runoff. Regression equation reveals that between the temperature, precipitation and runoff during 1960–2010 exhibited significant positive correlation (Fig. 4a, b), while there was a significant negative correlation between PET and runoff (Fig. 4c). In addition,  $F$  test and correlation significance test are conducted on 51 (or 50) samples in temperature, PET (or precipitation), and runoff. Confidence is taken as 99 % and  $P < 0.01$ . The 1 % significance level of the statistical test is passed, indicating that the linear regression coefficient can well display their relationship.

Based on Eq. (11) and Fig. 4, the elasticity of runoff in relation to temperature ( $\varepsilon_T$ ) and precipitation ( $\varepsilon_P$ ) are 0.1656 and 0.2078, respectively, which means that 10 % increase in temperature and precipitation will result in a 1.656 and 2.078 % increase in runoff. Meanwhile, the PET elasticity of runoff is estimated to be  $-1.1865$  ( $\varepsilon_{\text{PET}}$ ), which means that 1 % reduction in PET will lead to a 1.1865 % increase in runoff.

### 3.3 Impacts of CF on runoff

Based on CF change rates and elasticity of runoff in relation to precipitation, temperature, and PET, using Eq. (12), we can quantitatively evaluate the effects of climate factors on runoff change (Table 1). The results show that temperature change



**Fig. 2** Trend of runoff (a), temperature (b), precipitation (c), and PET (d) for Aksu River from 1960 to 2010


has the most importance effect on runoff change in recent 50 years, which may be related to runoff recharge larger proportions from glacier and snow melt water; on the other hand, the air temperature rises faster than other areas. Meanwhile, we can calculate the contributions of temperature, precipitation, and PET to runoff change in Aksu River with 45, 22, and 27 %, respectively, which indicates that the importance of runoff change for other factors (such as human activities, etc.) is about 6 %.

In addition, the research results in Table 1 are basically consistent with those Zheng et al. (2009) who came up with the other method obtained (Eq. 13), which implies that the suitability of this method (elasticity theory) to the area is credible and the conclusion of this article has the rationality


and reliability. The conclusion in two methods is that the importance of runoff change in Aksu River for climate factors is more than 90 %. Similarly, the research results of Chen et al. (2013) revealed that due to climate change, the increasing percentage in runoff of Kaidu River accounted for 90.5 % for the Tarim Basin, which is basically consistent with our results. Kaidu River also belongs to the Tarim Basin, plus, its runoff recharge proportion is from glacier and snow smelt water is also large. Thus, we can see that our results are believable.

#### 4 Discussion and conclusion

It is important to determine the impact of climate factors on runoff variation based on observation data. As a result, many methods have been widely adopted to assess the impact. However, it is hard to quantify the effects of temperature rising on runoff change. In this study, we defined a conceptual framework and applied the improved elasticity method to calculate the elasticity of runoff change to climate factors (CF, especially the temperature) in the past 50 years and then evaluated the contributions of CF to runoff change roughly. The results show that this method can be effective to evaluate quantitatively the influence of climate factors on runoff change. Concurrently, we realize that there are the uncertainties associated with estimating the impact of climate change on runoff using the elasticity method. First, based on



**Fig. 3** The step change point of annual runoff for Aksu River in recent 50 years



**Fig. 4 a–c** The regression relationships between the proportional change of runoff and proportional change of temperature, precipitation (remove an abnormal point,  $n=50$ ), and PET for Aksu River from 1960 to 2010

the assumption that every factor is independent of the others, the framework is used to estimate the proportional contribution of climate and other factors changes to runoff. However, in reality, the climate factors and other factors interact with each other. Second, a nonparametric statistical estimator of elasticity ( $\varepsilon$ ) was employed to identify the relationship between runoff and CF, which may involve inevitable uncertainty, for instance, the observation values may be influenced by undetermined factors. Third, we should attempt to use physics-based hydrological models to quantitatively evaluate the effects of climate factors on runoff change and verify it further in the future. All presented uncertainties would affect the results to some degree, and so estimation uncertainties need be further explored in further studies.

The contribution of climate factors (especially the temperature) to runoff change has not quantitatively been assessed

clearly for the arid region during the period 1960–2010. From the current study, we can conclude that the temperature change in Aksu River is the most important (45 %) to runoff variation, while the non-climate factors (such as human activities, etc.) are responsible for about 6 % of the runoff change in recent 50 years. Similarly, Chen et al. (2013) research results showed that due to human activities, the increasing percentage in runoff of Kaidu River in Tarim Basin only accounted for 9.5 % in the arid region, which is basically consistent with our results. Meanwhile, Huo et al. (2008) studied the effects of climate changes and water-related human activities on annual stream flows of the Shiyang River Basin in arid northwest China and pointed out that climate change was responsible for a large proportion of the runoff decreases in the upstream section of the catchment during the 1980s and 1990s, while human activities gave rise to runoff decreases in the

**Table 1** Quantitatively evaluating the effects of climate factors on runoff change on the basis of different climate elasticity estimators

Item	Change rate (%)	$\varepsilon_i$		CF result in runoff change ( $10^8 \text{ m}^3$ )		Importance of runoff change for CF (%)	
		This study	Equation (13)	This study	Equation (13)	This study	Equation (13)
Temperature	60 %	0.1656	0.1815	7.2	7.9	45 %	50 %
Precipitation	23 %	0.2078	0.1602	3.5	2.7	22 %	17 %
PET	5 %	-1.1865	-1.0507	4.3	3.8	27 %	24 %
Total	–	–	–	15	14.4	94 %	91 %

downstream for the same period. Our study area is the upstream of Aksu River, so regional climate is the important reason for the annual runoff change. Ma et al. (2008a) found that precipitation variability was the most important factor that decreased runoff for Zamu catchment in the arid region, which was different from our study. Because the Zamu Basin runoff recharge from mountain precipitation is the largest, while less runoff recharge proportion (1.4 %) is from glaciers melt water (Ma et al. 2008b). There are three reasons: (1) large runoff recharges proportion (about 42 %) from glaciers and snow melt water in Aksu River (Jiang et al. 2005). (2) The air temperature in the study area had a significant rising trend (Chen et al. 2009), at a rate of 0.28 °C/decade, higher than the average of entire globe (0.13 °C/decade) for the same period (IPCC 2007). (3) The selected hydrological station located in the mountain pass has a good representativeness in runoff change, while human activity intensity is weak in the mountains area.

**Acknowledgments** The research is supported by the National Basic Research Program of China (973 Program: 2010CB951003), Startup Fund Scientific Research Project of Qufu Normal University (BSQD20130102), National Natural Science Foundation of China (41171165, 41361093), Beijing Municipal Colleges and Universities High-level Talent Introduction and Training Program (IDHT20130322), and Funding Project for Academic Human Resources Development in Beijing Union University (BPHR2012E01). We are grateful to Dr. Hartmut Graßl and anonymous reviewers for their helpful comments on improving the manuscript.

## References

- Allen RG, Pereira LS, Raes D, Smith M (1998) Crop evapotranspiration: guidelines for computing crop water requirements. FAO Irrigation and Drainage Paper No. 56. FAO, Rome, Italy
- Chen Z, Chen Y (2014) Effects of climate fluctuations on runoff in the headwater region of the Kaidu River in northwestern China. *Front Earth Sci*. doi:10.1007/s11707-014-0406-2
- Chen YN, Xu ZX (2005) Plausible impact of global climate change on water resources in the Tarim River Basin. *Sci China Ser D Earth Sci* 48:65–73
- Chen YN, Xu CC, Hao XM, Li WH et al (2009) Fifty-year climate change and its effect on annual runoff in the Tarim River Basin, China. *Quatern Int* 208(1):53–61
- Chen ZS, Chen YN, Li WH (2012) Response of runoff to change of atmospheric 0°C level height in summer in arid region of Northwest China. *Sci China Ser D Earth Sci* 55:1533–1544
- Chen Z, Chen Y, Li B (2013) Quantifying the effects of climate variability and human activities on runoff for Kaidu River Basin in arid region of northwest China. *Theor Appl Climatol* 111:537–545
- Huo Z, Feng S, Kang S, Li W et al (2008) Effect of climate changes and water-related human activities on annual stream flows of the Shiyang River Basin in arid north-west China. *Hydrol Process* 22(16):3155–3167
- Intergovernmental Panel on Climate Change (IPCC) (2007) Climate change 2007: the physical science basis. Contribution of Working Group I to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change. In: Solomon S, et al. (ed). Cambridge Univ. Press, Cambridge, U. K., 996 pp
- Jiang Y, Zhou CH, Cheng WM (2005) Analysis on runoff supply and variation characteristics of Aksu drainage basin (In Chinese with English abstract). *J Nat Resour* 20(1):27–34
- Jiang SH, Ren LL, Yong B, Singh VP et al (2011) Quantifying the effects of climate variability and human activities on runoff from the Laohahe Basin in northern China using three different methods. *Hydrol Process* 25(16):2492–2505
- Jiang SH, Ren LL, Yong B, Fu C et al (2012) Analyzing the effects of climate variability and human activities on runoff from the Laohahe Basin in northern China. *Hydrol Res* 43(1–2):3–13
- Jin DL, Wu LB (2002) The application of elastic theory in economics. *Tech Econ* 172 (In Chinese)
- Kendall MG (1975) Rank correlation methods. Griffin, London
- Li H, Jiang Z, Liu X, Yang Q (2008) The relationship between the North Atlantic Oscillation and runoff variation of Aksu River in Xinjiang, China. *Acta Geograph Sin* 63(5):491–501
- Li BF, Chen YN, Shi X (2012a) Why does the temperature rise faster in the arid region of northwest China? *J Geophys Res* 117, D16115. doi:10.1029/2012JD017953
- Li BF, Chen YN, Chen ZS, Li WH (2012b) Trends in runoff versus climate change in typical rivers in the arid region of northwest China. *Quatern Int* 282:87–95
- Li BF, Chen YN, Shi X, Chen ZS (2013a) Temperature and precipitation changes in different environments in the arid region of northwest China. *Theor Appl Climatol* 112:589–596
- Li BF, Chen YN, Shi X, Chen ZS et al (2013b) Variations of temperature and precipitation of snowmelt period and its effect on runoff in the mountainous areas of northwest China. *J Geogr Sci* 23(1):17–30
- Li B, Su HB, Chen F, Li HB et al (2014a) Separation of the impact of climate change and human activity on streamflow in the upper and middle reaches of the Taoer River, northeastern China. *Theor Appl Climatol* 118:271–283
- Li F, Zhang G, Xu YJ (2014b) Separating the impacts of climate variation and human activities on runoff in the Songhua River Basin, northeast China. *Water* 6(11):3320–3338
- Liu DD, Chen XH, Lian YQ, Lou ZH (2010) Impacts of climate change and human activities on surface runoff in the Dongjiang River Basin of China. *Hydrol Process* 24:1487–1495
- Ma Z, Kang S, Zhang L, Tong L, Su X (2008a) Analysis of impacts of climate change and human activity on streamflow for a river basin in arid region of northwest China. *J Hydrol* 352:239–249
- Ma G, Liu J, Lin D, Chen N (2008b) Status of water use and its Eco2environmental effects in Shiyanghe River Basin (in Chinese with English abstract). *J Desert Res* 28(3):592–597
- Mann HB (1945) Nonparametric tests against trend. *Econometrica* 13: 124–259
- Meng D, Mo X (2012) Assessing the effect of climate change on mean annual runoff in the Songhua River Basin, China. *Hydrol Process* 26(7):1050–1061
- Milly PCD, Dunne KA, Vecchia AV (2005) Global pattern of trends in streamflow and water availability in a changing climate. *Nature* 438: 347–350
- Ren G, Xu M, Chu Z, Guo J et al (2005) Changes of surface air temperature in China during 1951–2004 (in Chinese with English abstract). *Clim Environ Res* 10(4):717–727
- Sankarasubramanian A, Vogel RM, Limburner JF (2001) Climate elasticity of streamflow in the United States. *Water Resour Res* 37: 1771–1781. doi:10.1029/2000WR900330
- Schaake JC (1990) From climate to flow. In: Waggoner PE (ed) Climate change and U.S. water resources, chap. 8. John Wiley, New York. pp. 177–206
- Sneyers R (1975) Sur l'analyse statistique des séries d'observations. O.M.M., Note Technique No 143, Gencve, Suisse



- Song XM, Zhang JY, Zhan CS, Liu CZ (2013) Review for impacts of climate change and human activities on water cycle (in Chinese with English abstract). *J Hydraul Eng* 44(7):779–790
- Wang S, Yan Y, Yan M, Zhao X (2012) Quantitative estimation of the impact of precipitation and human activities on runoff change of the Huangfuchuan River Basin. *J Geogr Sci* 22(5):906–918
- Wang W, Shao Q, Yang T, Peng S et al (2013) Quantitative assessment of the impact of climate variability and human activities on runoff changes: a case study in four catchments of the Haihe River Basin, China. *Hydrol Process* 27:1158–1174
- Xu JH, Chen YN, Lu F, Li WH et al (2011) The Nonlinear trend of runoff and its response to climate change in the Aksu River, western China. *Int J Climatol* 31(5):687–695
- Yang X, Yan J, Liu B (2005) The analysis on the change characteristic and driving forces of Wuding River runoff (in Chinese with English abstract). *Adv Earth Sci* 20(6):637–642
- Ye X, Zhang Q, Liu J, Li X et al (2009) Impacts of climate change and human activities on runoff of Poyang Lake Catchment (in Chinese with English abstract). *J Glaciol Geocryol* 31(5):835–842
- Ye X, Zhang Q, Liu J, Li X et al (2013) Distinguishing the relative impacts of climate change and human activities on variation of streamflow in the Poyang Lake catchment, China. *J Hydrol* 494:83–95
- You QL, Kang SC, Aguilar E et al (2011) Changes in daily climate extremes in China and their connection to the large scale atmospheric circulation during 1961–2003. *Clim Dyn* 36(11–12):2399–2417
- Yu PJ, Xu HL, Liu SW, An HY et al (2011) The nonlinear characteristics of annual runoff change in Aksu River. *J Nat Resour* 26(8):1412–1422
- Zang C, Liu J, Jiang L, Gerten D (2013) Impacts of human activities and climate variability on green and blue water flows in the Heihe River Basin in northwest China. *Hydrol Earth Syst Sci Discuss* 10:9477–9504
- Zeng S, Xia J, Du H (2014) Separating the effects of climate change and human activities on runoff over different time scales in the Zhang River Basin. *Stoch Environ Res Risk Assess* 28(2):401–413
- Zhang XQ, Sun Y, Mao WF, Liu YY, Ren Y (2010) Regional response of temperature change in the arid regions of China to global warming (in Chinese with English abstract). *Arid Zone Res* 27(4):592–599
- Zhang Q, Li JF, Chen YD, Chen XH (2011) Observed changes of temperature extremes during 1960–2005 in China: natural or human-induced variations? *Theor Appl Climatol* 106:417–431
- Zhang AJ, Zhang C, Fu GB et al (2012) Assessments of impacts of climate change and human activities on runoff with SWAT for the Huifa River Basin, northeast China. *Water Resour Manag* 26(8):2199–2217
- Zheng HX, Zhang L, Zhu RR, Liu CM et al (2009) Responses of streamflow to climate and land surface change in the headwaters of the Yellow River Basin. *Water Resour Res* 45, W00A19. doi:10.1029/2007WR-006665